



A recent update on the connection between dietary phytochemicals and skin cancer: emerging understanding of the molecular mechanism

Harpreet Singh, PhD^{a,*}, Arun Kumar Mishra, PhD^b, Sourav Mohanto, PhD^c, Arvind Kumar, PhD^a, Amrita Mishra, PhD^d, Ruhul Amin, PhD^{e,*}, Chellappan Ronald Darwin, PhD^f, Talha Bin Emran, PhD^{g,*}

Abstract

Constant exposure to harmful substances from both inside and outside the body can mess up the body's natural ways of keeping itself in balance. This can cause severe skin damage, including basal cell carcinoma (BCC), squamous cell carcinoma (SCC), and melanoma. However, plant-derived compounds found in fruits and vegetables have been shown to protect against skin cancer-causing free radicals and other harmful substances. It has been determined that these dietary phytochemicals are effective in preventing skin cancer and are widely available, inexpensive, and well-tolerated. Studies have shown that these phytochemicals possess anti-inflammatory, antioxidant, and antiangiogenic properties that can aid in the prevention of skin cancers. In addition, they influence crucial cellular processes such as angiogenesis and cell cycle control, which can halt the progression of skin cancer. The present paper discusses the benefits of specific dietary phytochemicals found in fruits and vegetables, as well as the signaling pathways they regulate, the molecular mechanisms involved in the prevention of skin cancer, and their drawbacks.

Keywords: antioxidant, basal cell carcinoma, dietary phytochemicals, melanoma skin cancer, squamous cell carcinoma

Introduction

Skin structure

The skin is the largest organ in the human body and serves as a barrier between the internal and external environment. It plays a

HIGHLIGHTS

- Skin cancer is a type of cancer that arises from the uncontrolled growth of cells in the skin. There are three main types of skin cancer.
- This mutation also leads to the activation of the MAPK/ERK signaling pathway and abnormally promotes cell growth and division.
- The BRAF and NRAS genes are members of the RAS family of genes, which regulate cell growth and division.
- The CDKN2A gene encodes two proteins, p16INK4a and p14ARF, which act as tumor suppressors by regulating cell growth and division.
- Signaling pathways (MAPK pathway, PI3K/Akt pathway, Wnt pathway, Hedgehog pathway, Notch pathway, etc.) are involved in cell growth and proliferation.
- Phytochemicals found in fruits and vegetables, as well as the signaling pathways they regulate, the molecular mechanisms involved in the prevention of skin cancer.

^aSchool of Pharmaceutical Sciences, IFTM University, Moradabad, Uttar Pradesh,

^bSOS School of Pharmacy, IFTM University, Moradabad, Uttar Pradesh,

^cDepartment of Pharmaceutics, Yenepoya Pharmacy College & Research Centre, Yenepoya (Deemed to be University), Mangalore, Karnataka, ^dSchool of Pharmaceutical Sciences, Delhi Pharmaceutical Sciences and Research University, New Delhi, ^eFaculty of Pharmaceutical Science, Assam downtown University, Panikhaiti, Gandhinagar, Guwahati, Assam, ^fHOD, Department of Pharmacology, School of Pharmaceutical Sciences, VISTAS, Chennai, India and ^gDepartment of Pharmacy, Faculty of Allied Health Sciences, Daffodil International University, Dhaka, Bangladesh

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

*Corresponding authors. Address: School of Pharmaceutical Sciences, IFTM University, Moradabad 244102, Uttar Pradesh, India. Tel.: +91 88659 34783. E-mail: harpreetproctor@rediffmail.com (H. Singh); Faculty of Pharmaceutical Science, Assam downtown University, Panikhaiti, Gandhinagar, Guwahati 781026, Assam, India. Tel.: +91 7002 942887. E-mail: ruhulgp18@gmail.com (R. Amin); Department of Pharmacy, Faculty of Allied Health Sciences Daffodil International University, Dhaka 1207, Bangladesh. Tel.: +88 018 199 42214. E-mail: talhabmb@bgctub.ac.bd (T.B. Emran).

Copyright © 2024 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Annals of Medicine & Surgery (2024) 86:5877–5913

Received 27 May 2024; Accepted 8 July 2024

Published online 7 August 2024

<http://dx.doi.org/10.1097/MS9.0000000000002392>

crucial role in protecting the body from physical, chemical, and biological damage, as well as in regulating body temperature, maintaining hydration, and providing sensory input^[1]. The skin is composed of three main layers: the epidermis, dermis, and hypodermis (subcutaneous tissue) (Fig. 1)^[2]. Each layer has its own unique structure and function, which work together to provide a protective and functional barrier. The epidermis is the outermost layer of the skin and is composed of several layers of keratinized epithelial cells. The thickness of the epidermis varies depending on the location on the body, with the thickest epidermis found on the soles of the feet and the thinnest on the eyelids. The epidermis is responsible for protecting the body from




Figure 1. Skin anatomy (created with BioRender).

physical damage, UV radiation, and infection^[3]. The dermis is the second layer of the skin and is located beneath the epidermis. The dermis is composed of connective tissue and is responsible for providing support and elasticity to the skin. The dermis also contains blood vessels, nerves, and sensory receptors. The hypodermis, also known as the subcutaneous tissue, is the deepest layer of the skin. The hypodermis is composed of adipose tissue and connective tissue and serves as a cushion for the underlying muscles and bones^[4].

There are several types of cells present in the skin, each with its own unique function. Some of the main cell types in the skin include Keratinocytes, Melanocytes, Langerhans cells, Merkel cells, and Fibroblasts^[5]. Skin cancer is the most common type of cancer in the United States and is caused by the uncontrolled growth of abnormal cells in the skin. The different cell types in the skin play a critical role in protecting against skin cancer. For example, melanocytes produce melanin, which helps to protect against UV radiation and reduces the risk of skin cancer. Langerhans cells are involved in the immune response and can help to identify and destroy abnormal cells before they can develop into skin cancer. Additionally, fibroblasts produce the extracellular matrix, which helps to prevent the uncontrolled growth and spread of abnormal cells. Overall, the different types of cells in the skin work together to maintain the integrity of the skin and protect against the development of skin cancer. Regular skin checks and sun protection can also help to reduce the risk of skin cancer^[6,7].

Molecular mechanism of skin cancer

Skin cancer is a type of cancer that arises from the uncontrolled growth of cells in the skin. There are three main types of skin cancer: basal cell carcinoma (BCC), squamous cell carcinoma (SCC), and melanoma^[8–10]. The molecular mechanisms underlying skin cancer development involve mutations in several

genes and pathways that regulate cell growth, division, and apoptosis^[11,12]. BCC is the most common type of skin cancer, accounting for ~80% of all cases. It typically develops in the basal cells, which are located in the deepest layer of the epidermis. The molecular mechanisms underlying BCC development involve mutations in the sonic hedgehog (SHH) signaling pathway. The SHH signaling pathway plays an important role in embryonic development, and its activation is necessary for the development of the skin, hair, and nails. In adults, the SHH pathway is normally inactive in the skin. However, mutations in genes that regulate the SHH pathway can lead to its activation and the development of BCC^[13,14]. The most common mutation in BCC is in the Patched-1 (PTCH1) gene, which normally acts as a tumor suppressor by inhibiting the SHH pathway. Mutations in PTCH1 lead to the activation of the SHH pathway, which results in the uncontrolled growth of basal cells and the development of tumors^[15,16]. Other genetic and environmental factors can also contribute to the development of BCC. Mutations in the Tumor Protein 53 (TP53) gene, which is another tumor suppressor gene, can also lead to BCC development. Environmental factors, such as exposure to arsenic and ionizing radiation, can also increase the risk of developing BCC^[17,18]. SCC is the second most common type of skin cancer, accounting for ~16% of all cases. It typically develops in the squamous cells, which are located in the upper layers of the epidermis^[19]. The molecular mechanisms underlying SCC development involve mutations in several genes, including TP53, Cyclin-dependent kinase inhibitor 2A (CDKN2A), and Harvey rat sarcoma viral oncogene homolog (HRAS). Like BCC, mutations in TP53 and CDKN2A are commonly found in SCC. TP53 mutations can lead to the uncontrolled growth of squamous cells and the development of tumors^[20,21]. CDKN2A mutations can also contribute to SCC development by preventing cells from undergoing apoptosis, which is a process that helps to eliminate damaged cells from the body^[22,23]. Mutations in the HRAS gene, which is a member of

the Rat Sarcoma (RAS) family of genes, are also found in SCC. The RAS genes normally regulate cell growth and division, but mutations in these genes can lead to the development of tumors. In SCC, mutations in HRAS lead to the uncontrolled growth of squamous cells and the development of tumors^[24,25]. Melanoma is a type of skin cancer that arises from the uncontrolled growth of melanocytes, which are the cells that produce the pigment melanin. Melanoma is the most dangerous type of skin cancer, as it can metastasize and spread to other parts of the body. The molecular mechanisms underlying melanoma development also involve mutations in several genes and pathways that regulate cell growth, division, and apoptosis^[26–29].

B-Raf proto-oncogene, serine/threonine kinase (BRAF), and neuroblastoma RAS viral oncogene homolog (NRAS) mutations

The BRAF and NRAS genes are members of the RAS family of genes, which regulate cell growth and division. Mutations in BRAF and NRAS are commonly found in melanoma and can lead to the uncontrolled growth of melanocytes and the development of tumors^[27]. The BRAF gene is mutated in ~50% of all melanomas. The most common mutation is a substitution of valine for glutamic acid at position 600 (BRAFV600E). This mutation leads to the activation of the mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/ERK) signaling pathway, which promotes cell growth and division. The NRAS gene is mutated in ~15% of all melanomas. The most common mutation is a substitution of glutamine for arginine at position 61 (NRASQ61R). This mutation also leads to the activation of the MAPK/ERK signaling pathway and promotes cell growth and division^[30,31].

CDKN2A mutations

The CDKN2A gene encodes two proteins, p16INK4a and p14ARF, which act as tumor suppressors by regulating cell growth and division. Mutations in CDKN2A are commonly found in melanoma and can lead to the uncontrolled growth of melanocytes and the development of tumors. Loss of p16INK4a function is found in ~50% of all melanomas. This loss can occur through mutations in the CDKN2A gene or through epigenetic silencing of the gene. Loss of p16INK4a function leads to the activation of the cyclin-dependent kinase (CDK) 4/6 pathway, which promotes cell growth and division. Loss of p14ARF function is found in ~20% of all melanomas. This loss can occur through mutations in the CDKN2A gene or through epigenetic silencing of the gene. Loss of p14ARF function leads to the stabilization of the oncoprotein mouse double minute 2 (MDM2) homolog, which promotes the degradation of the tumor suppressor protein p53^[32,33].

TP53 mutations

The TP53 gene encodes the tumor suppressor protein p53, which regulates cell growth, division, and apoptosis. Mutations in TP53 are found in ~25% of all melanomas and can lead to the uncontrolled growth of melanocytes and the development of tumors^[34–36]. Loss of p53 function can occur through mutations in the TP53 gene or through the stabilization of the oncoprotein MDM2, which promotes the degradation of p53. Loss of p53

function leads to the inhibition of apoptosis and the promotion of cell growth and division^[37,38].

Skin cancer therapy via diet-based phytochemicals

Dietary phytochemicals are naturally occurring compounds found in plant-based foods, including fruits, vegetables, whole grains, nuts, seeds, and herbs. Phytochemicals are not considered essential nutrients, but they have been shown to have a wide range of health benefits, including antioxidant, anti-inflammatory, anticancer, and neuroprotective properties^[39–42]. There are thousands of different phytochemicals, each with its unique chemical structure and biological activity. Some of the most well-known and studied phytochemicals include flavonoids, carotenoids, phenolic acids, and lignans^[43–46]. Flavonoids, such as quercetin and kaempferol, are potent antioxidants that have been shown to have anti-inflammatory and anticancer effects^[47,48]. Carotenoids, such as β -carotene, are responsible for the red, orange, and yellow colors in fruits and vegetables and have been shown to have potent antioxidant and anticancer properties^[49,50]. Phenolic acids, such as caffeic acid and ferulic acid, are found in a wide range of plant-based foods and have been shown to have antioxidant and anti-inflammatory effects^[43]. Lignans are phytoestrogens found in flaxseeds, sesame seeds, and whole grains and have been shown to have anticancer effects^[51,52]. The health benefits of dietary phytochemicals are believed to come from their ability to interact with cellular signaling pathways, enzymes, and other molecules in the body, thereby modulating various biological processes (Table 1A–N)^[127–129].

Overall, skin cancer therapy via diet-based phytochemicals is a promising approach for the prevention and treatment of skin cancer. Further research is needed to identify the optimal dosage, duration, and combination of phytochemicals for maximum therapeutic efficacy. However, the evidence to date supports the inclusion of phytochemical-rich foods in the diet as an important strategy for skin cancer prevention and therapy. The goal of this review article is to give a basic understanding of phytochemicals in the context of diet by describing their most important sources, chemical classes, and ability to prevent skin cancer.

Rosmarinic acid

Rosmarinic acid is a naturally occurring polyphenol found in various plant species, particularly in the *Lamiaceae* family^[130,131]. Some of the most common botanical sources of rosmarinic acid include Rosemary (*Rosmarinus officinalis*), Sage (*Salvia officinalis*), Lemon balm (*Melissa officinalis*), Oregano (*Origanum vulgare*), Thyme (*Thymus vulgaris*), Mint (*Mentha* spp.), Basil (*Ocimum basilicum*), Perilla (*Perilla frutescens*), Lavender (*Lavandula angustifolia*), and Peppermint (*Mentha piperita*)^[132–134].

Rosmarinic acid has been shown to have potential chemopreventive effects against skin cancer through multiple molecular mechanisms^[135]. Rosmarinic acid has potent antioxidant properties, which can protect skin cells from oxidative stress caused by environmental factors such as UV radiation. This can prevent DNA damage and mutations that can lead to the development of skin cancer^[136–141]. One mechanism through which rosmarinic acid may prevent skin cancer is by inhibiting the activity of enzymes called matrix metalloproteinases (MMPs). MMPs play a crucial role in the breakdown of collagen and other extracellular matrix components. Overactive MMPs can lead to the destruction

Table 1
Some important dietary phytochemicals effective in preventing skin cancer.








S. no.	Dietary phytochemical	Structure	IUPAC name	Signaling pathway	References
A					
1.	Rosmarinic acid		(2S)-2-[(3-(3,4-dihydroxyphenyl)-l-alanyloxy]-3-(4-hydroxyphenyl)propanoic acid	(1) Inhibition of PI3K/AKT/mTOR pathway. (2) Inhibition of MAPK/ERK pathway. (3) Inhibit the activation of NF-κB. (4) Inhibit the Wnt/β-catenin pathway by decreasing the expression of β-catenin and its downstream target genes, leading to decreased cell proliferation and invasion.	[53] [54] [55] [56,57]
B					
2.	Allicin		2-propene-1-sulfinothioic acid S-2-propen-1-yl ester	(1) Inhibit the activation of NF-κB, potentially reducing the risk of various types of cancer development, including skin cancer. (2) Inhibit the activation of the Wnt/β-catenin pathway, potentially reducing the risk of various types of cancer development, including skin cancer.	[58] [59]
C					
3.	Sulforaphane		(RS)-1-isothiocyanato-4-(methylsulfinyl)butane	(1) Activate the Nrf2–Keap1 pathway, leading to increased production of antioxidant and detoxification enzymes that help protect against the damaging effects of environmental toxins and UV radiation. (2) Inhibit the activation of these PI3K/Akt and MAPK pathways potentially reducing the risk of cancer development, including skin cancer.	[60] [61,62]
D					
4.	Ellagic acid		2,3,7,8-tetrahydroxychromeno [5,4,3-cde]chromene-5,10-dione	(1) Inhibit the activation of NF-κB signaling pathway to prevent the proliferation of skin cancer cells and induce apoptosis. (2) Activate the p53 signaling pathway, which can induce cell cycle arrest and apoptosis in skin cancer cells. (3) Modulation of MAPK signaling pathway. (4) Inhibition of PI3K/Akt signaling pathway.	[63–65] [66] [63] [67]
E					
5.	Betulinic acid		(1R,3aS,5aR,5bR,7aR,9S,11aR,11bR,13aR,13bR)-9-hydroxy-5a,5b,8,8,11a-pentamethyl-1-prop-1-en-2-yl-1,2,3,4,5,6,7,7a,9,10,11,11b,12,13,13a,13b-hexadecahydrocyclopenta[a]chrysene-3a-carboxylic acid	(1) Inhibit the activation of NF-κB signaling pathway. (2) Activation of p53 signaling pathway. (3) Modulation of MAPK signaling pathway. (4) Inhibition of PI3K/Akt signaling pathway. (5) Inhibition of Notch signaling pathway.	[68] [69,70] [71] [72] [73]
F					
6.	Apigenin		4',5,7-trihydroxyflavone	(1) Inhibition of PI3K/Akt signaling pathway. (2) Inhibition of MAPK signaling pathway. (3) By inhibiting the activation of signal transducer and activator of transcription 3 (STAT3). (4) Apigenin can activate the p53 signaling pathway, which can induce cell cycle arrest and apoptosis in skin cancer cells. (5) By inhibiting NF-κB signaling pathway.	[74] [75] [76] [77] [78,79]
G					
7.	Gingerol		(S)-5-hydroxy-1-(4-hydroxy-3-methoxyphenyl)-3-decanone	(1) Inhibition of PI3K/Akt/mTOR signaling pathway. (2) Inhibition of NF-κB signaling pathway. (3) Activation of nuclear factor erythroid 2-related factor 2 (Nrf2) signaling pathway. (4) Inhibition of MAPK signaling pathway. (5) Induction of apoptosis.	[80] [81] [82] [83] [80]

Table 1
(Continued)









S. no.	Dietary phytochemical	Structure	IUPAC name	Signaling pathway	References
H					
8.	Quercetin		3,3',4',5,7-pentahydroxyflavone	(1) Inhibition of oxidative stress. (2) Inhibit the production of pro-inflammatory cytokines such as interleukin-1 β (IL-1 β), interleukin-6 (IL-6), and tumor necrosis factor- α (TNF- α). (3) Inhibition of cell proliferation. (4) Inhibition of angiogenesis.	[84] [85,86] [58] [87]
I					
9.	Kaempferol		3,5,7-trihydroxy-2-(4-hydroxyphenyl)-4H-1-benzopyran-4-one	(1) Inhibition of oxidative stress. (2) Inhibition of inflammation. (3) Inhibition of cell proliferation. (4) Inhibition of angiogenesis.	[88,89] [90] [88] [91,92]
10.	Resveratrol		3,5,4'-trihydroxy-trans-stilbene	(1) Inhibition of oxidative stress. (2) Inhibition of inflammation. (3) Inhibition of cell proliferation. (4) Inhibition of angiogenesis.	[93] [94,95] [96] [97,98]
J					
11.	Curcumin		(1E,6E)-1,7-bis(4-hydroxy-3-methoxyphenyl)hepta-1,6-diene-3,5-dione	(1) Inhibition of oxidative stress. (2) Inhibition of inflammation. (3) Inhibition of cell proliferation. (4) Inhibition of angiogenesis.	[99,100] [101] [102,103] [104,105]
K					
12.	Epigallocatechin gallate		[(2R,3R)-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)-3,4-dihydro-2H-1-benzopyran-3-yl] 3,4,5-trihydroxybenzoate	(1) Inhibition of oxidative stress. (2) Inhibition of inflammation. (3) Inhibition of cell proliferation. (4) Inhibition of angiogenesis.	[106,107] [108,109] [110,111] [112,113]
L					
13.	β -Carotene		1,3,3-trimethyl-2-[3,7,12,16-tetramethyl-18-(2,6,6-trimethyl-1-cyclohexenyl)octadeca-1,3,5,7,9,11,13,15,17-nonaenyl]cyclohexene	(1) Inhibition of oxidative stress. (2) Modulation of immune function. (3) Inhibition of inflammation. (4) Regulation of cell cycle and apoptosis. (5) Inhibition of angiogenesis.	[114,115] [116,117] [118,119] [115] [120]
M					
14.	Caffeic acid		3-(3,4-dihydroxyphenyl)prop-2-enoic acid	(1) Inhibit the activation of NF- κ B by preventing the degradation of its inhibitor protein, I κ B α . (2) Regulates the expression of mitogen-activated protein kinases (MAPKs), which regulate cell proliferation and survival, and cyclooxygenase-2 (COX-2), which promotes inflammation and cell proliferation.	[121,122] [123,124]

Table 1

(Continued)

S. no.	Dietary phytochemical	Structure	IUPAC name	Signaling pathway	References
N					
15.	Ferulic acid		(E)-3-(4-hydroxy-3-methoxyphenyl)prop-2-enoic acid	(1) Inhibit the activation of NF-κB by preventing the degradation of its inhibitor protein, IκBα. (2) Regulates the expression of MAPKs, which regulate cell proliferation and survival, and COX-2, which promotes inflammation and cell proliferation.	[125] [126]

of the extracellular matrix, which can promote the development of skin cancer. Rosmarinic acid has been shown to inhibit the activity of MMPs, which may help prevent skin cancer^[142,143]. Chronic inflammation is a key contributor to skin cancer development. Rosmarinic acid has been shown to have anti-inflammatory effects by inhibiting the production of pro-inflammatory cytokines and enzymes such as cyclooxygenase-2 (COX-2) and inducible nitric oxide synthase (iNOS). By reducing inflammation, rosmarinic acid can prevent the progression of precancerous lesions to skin cancer^[144,145]. Rosmarinic acid has been shown to induce apoptosis (programmed cell death) in skin cancer cells, which can inhibit their growth and proliferation^[146–150]. Rosmarinic acid has been shown to arrest the cell cycle of skin cancer cells, preventing them from dividing and proliferating^[151,152]. Angiogenesis, the formation of new blood vessels, is necessary for tumor growth and progression. Rosmarinic acid has been shown to inhibit angiogenesis in skin cancer cells, preventing their growth and spread^[153].

In recent years, researchers have investigated the molecular mechanisms underlying the effects of rosmarinic acid on skin cancer cells, including its ability to induce apoptosis, inhibit inflammation, and regulate cell cycle progression. For example, Gupta *et al.* investigated the toxic effects of UVB radiation on the skin and the potential therapeutic effects of plant-based natural agents. The study found that UVB exposure induced endoplasmic reticulum (ER) stress and inhibited mitophagy, leading to intracellular damage and apoptosis. Treatment with the natural agent rosmarinic acid prevented intracellular damage by alleviating ER stress and promoting mitophagy. The study highlights the potential of rosmarinic acid as a therapeutic agent for photodamage and provides mechanistic insights into the toxic effects of UVB radiation on skin^[154]. Another study, conducted by Huang *et al.*, investigated the anticancer effects of rosmarinic acid in melanoma cells by downregulating (a disintegrin and metalloproteinase 17) ADAM17. Results showed that rosmarinic acid treatment reduced cell viability, proliferation, migration, and invasion abilities while increasing apoptosis and reducing melanin content. Rosmarinic acid also inhibited the expression of ADAM17/epidermal growth factor receptor (EGFR)/protein kinase B (AKT)/glycogen synthase kinase 3 beta (GSK3β), which was further suppressed by TPD, an ADAM17 inhibitor. The study concludes that rosmarinic acid exerts an inhibitory effect on melanoma cell growth and promotes apoptosis, potentially through the inhibition of the ADAM17/EGFR/AKT/GSK3β axis^[155]. A study by Sharmila *et al.* investigated the mechanisms by which rosmarinic acid affects the expression of MAPK signaling proteins and their downstream targets in mice with dermal cancer, as well as to analyze the docking

interaction of rosmarinic acid with the extracellular signal-regulated kinase 2 (ERK2) protein. Dermal cancer was induced in mice by applying 7,12-dimethylbenz[a]anthracene (DMBA), and various analyses were performed to observe the expression of proteins related to MAPK signaling, as well as histopathological changes. The study found that rosmarinic acid significantly reduced the expression of various proteins related to MAPK signaling in dermal tissues and inhibited the activation of ERK2 protein, which may contribute to the inhibitory effect of rosmarinic acid on dermal cancer in mice^[156]. The study conducted by Lukmanul Hakkim *et al.* aimed to investigate whether natural antioxidants including caffeic acid, rosmarinic acid, trans-cinnamic acid, *p*-coumaric acid, and hydroxyphenyllactic acid could offer radiation protection for skin cells. Non-toxic concentrations of these compounds were tested for radiation protection in human keratinocytes. Results showed that pretreatment with caffeic acid, rosmarinic acid, and trans-cinnamic acid could protect skin cells by scavenging γ-radiation-induced reactive oxygen species and decreasing the number of post-irradiation DNA double-strand break foci. The inclusion of these compounds in chemo-radiotherapy could potentially facilitate achieving multiple target protection, including anticancer and skin radio protection^[157].

Rosmarinic acid has been shown to regulate several major signaling pathways in the human body including:

The phosphatidylinositol 3-kinase/protein kinase B/ mammalian target of rapamycin (PI3K/AKT/mTOR) pathway

The PI3K/AKT/mTOR pathway is frequently activated in various types of cancer, including skin cancer. Rosmarinic acid has been shown to inhibit this pathway by decreasing the phosphorylation of AKT and mTOR, leading to decreased cell proliferation and survival^[53].

The MAPK/ERK pathway

The MAPK/ERK pathway is another important signaling pathway involved in skin cancer development. Rosmarinic acid has been shown to inhibit this pathway by decreasing the phosphorylation of ERK, leading to decreased cell proliferation and invasion^[54].

The nuclear factor-κB (NF-κB) pathway

The NF-κB pathway is a transcription factor that plays a critical role in inflammation and cancer development. Rosmarinic acid has been shown to inhibit the activation of NF-κB by preventing the degradation of the inhibitor of kappa B alpha (IκBα), leading to decreased inflammation and cell proliferation^[55].

The Wnt/ β -catenin pathway

The Wnt/ β -catenin pathway is involved in the regulation of cell proliferation, differentiation, and apoptosis. Dysregulation of this pathway has been linked to various types of cancer, including skin cancer. Rosmarinic acid has been shown to inhibit this pathway by decreasing the expression of β -catenin and its downstream target genes, leading to decreased cell proliferation and invasion^[56,57].

The ability of rosmarinic acid to regulate major signaling pathways in the human body suggests that it may have therapeutic potential for the prevention and treatment of various diseases, including cancer^[158]. While rosmarinic acid has shown promising anticancer properties *in vitro* and in animal studies, there are several potential drawbacks and limitations to its use in treating skin cancer^[131,159]. While rosmarinic acid is generally considered safe, there are some potential side effects, including nausea, vomiting, and allergic reactions^[160]. Rosmarinic acid has poor bioavailability, which means that it may not be absorbed well by the body when taken orally or applied topically^[161]. The amount of rosmarinic acid in herbal supplements or preparations can vary widely, which makes it difficult to establish a standardized dose or formulation for treating skin cancer. This may limit its effectiveness in treating skin cancer^[162,163]. More research is needed to determine its safety and efficacy and to optimize the dosage and delivery of rosmarinic acid for maximum efficacy.

Allicin

Allicin is a natural compound found in garlic (*Allium sativum*)^[164]. Allicin belongs to the class of secondary metabolites called organosulfur compounds. These compounds are characterized by the presence of sulfur atoms in their chemical structure and are often produced by plants as a defense mechanism against pests and pathogens^[165–167]. When garlic is crushed or chopped, an enzyme called alliinase is activated, which converts alliin, a sulfur-containing amino acid derivative, into allicin^[168]. Allicin is responsible for the characteristic odor and flavor of fresh garlic, and it is also believed to be responsible for many of the health benefits associated with garlic consumption^[169,170]. The molecular mechanism of allicin's action against skin cancer is not completely understood, but several studies have suggested that it targets multiple pathways involved in cancer development and progression^[171]. A study investigated the effect of allicin on the migration and invasion of human melanoma cells (A375 and SK-MEL-28). Allicin was found to inhibit the migration and invasion of melanoma cells by downregulating the expression of genes involved in epithelial–mesenchymal transition (EMT), a process that allows cancer cells to acquire invasive and metastatic properties. Allicin was also found to downregulate the expression of the COX-2 gene, which is involved in inflammation and cell proliferation. Furthermore, allicin was found to inhibit the activation of the NF- κ B pathway and the expression of its downstream target genes, which are involved in cell survival, inflammation, and tumor progression^[58]. In a study conducted by Wang *et al.*, it was observed that several compounds, including allicin, allyl sulfides, ajoene, diallyl trisulfide (DATS), and S-allyl cysteine (SAC), have shown anticancer activity against various types of cancer, including skin cancer. DATS is more potent than mono- and disulfides against skin cancer. DATS inhibits cell growth of human melanoma A375 cells and basal cell carcinoma

(BCC) cells by increasing the levels of intracellular reactive oxygen species (ROS) and DNA damage, inducing G2/M arrest, endoplasmic reticulum (ER) stress, and mitochondria-mediated apoptosis, including the caspase-dependent and caspase-independent pathways^[171]. In one of the studies, Jobani *et al.* aimed to investigate the potential of allicin to sensitize malignant melanoma cells to all-trans retinoic acid (ATRA) therapy. The CD44+ and CD117+ melanoma cell subpopulations were sorted, and the effects of ATRA, allicin, and allicin/ATRA on cell proliferation and cell cycle arrest were examined. The results showed that CD44+ melanoma cells were more resistant to ATRA and allicin than CD117+ cells. However, allicin was found to sensitize melanoma cells to ATRA-induced cell death, and the combination treatment significantly reduced the IC₅₀ value obtained for ATRA alone in CD44+ melanoma cells. Furthermore, allicin and ATRA combination treatment showed inhibitory effects on CD44+ and CD117+ melanoma cells, and allicin alone reduced matrix metalloproteinase-9 (MMP-9) mRNA expression in both cell subpopulations. The findings suggest that allicin may reinforce the ATRA-mediated inhibitory effects on melanoma cells, providing a new approach for the treatment of malignant melanoma^[172].

In another study conducted by Omar and Al-Wabel, it was observed that garlic contains chemical compounds that have been shown to protect against several diseases, including cancer, particularly in the stomach, colorectal, breast, and skin. The protective effects are related to the presence of organosulfur compounds (like allicin, etc.), which inhibit carcinogenesis in various experimental animals. The compounds modulate the activity of several metabolizing enzymes and inhibit the formation of DNA adducts. Antiproliferative activity has been observed in tumor cell lines, possibly mediated by induction of apoptosis and alterations of the cell cycle. Garlic's organosulfur compounds are potential cancer-preventive agents, but clinical trials are necessary to define the effective dose with no toxicity in humans^[173].

Allicin has been shown to scavenge free radicals and protect against oxidative damage induced by UV radiation, which can contribute to the development of skin cancer^[174]. Allicin has been shown to suppress the production of inflammatory cytokines, such as interleukin-1 beta (IL-1 β) and tumor necrosis factor-alpha (TNF- α), which can promote the development and progression of skin cancer^[175]. Allicin has been shown to induce apoptosis (programmed cell death) in cancer cells, including skin cancer cells, by activating caspases and other apoptosis-related proteins^[176]. Allicin has been shown to arrest the cell cycle of cancer cells, preventing them from dividing and multiplying, which can help to prevent the growth and spread of skin cancer^[177]. Allicin has been shown to inhibit the formation of new blood vessels, which is a critical step in the growth and spread of cancer^[178].

The major signaling pathways that have been reported to be regulated by allicin are as follows:

NF- κ B pathway

Allicin has been shown to inhibit the activation of NF- κ B, a transcription factor that plays a key role in inflammation and cancer development^[58].

PI3K/Akt/mTOR pathway

Allicin has been shown to inhibit the activation of this pathway, which is frequently dysregulated in cancer cells and promotes cell survival, growth, and proliferation^[179].

Wnt/ β -catenin pathway

Allicin has been shown to inhibit the activation of this pathway, which is frequently dysregulated in cancer cells and promotes cell proliferation and survival^[59].

MAPK pathway

Allicin has been shown to inhibit the activation of the MAPK pathway, which plays a critical role in cell proliferation, differentiation, and survival^[180].

STAT3 pathway

Allicin has been shown to inhibit the activation of STAT3, a transcription factor that plays a critical role in inflammation and cancer development^[181].

Overall, these pathways play important roles in regulating cell growth, survival, and inflammation and dysregulation of these pathways can contribute to the development and progression of skin cancer. By regulating these pathways, allicin has the potential to prevent or slow down the progression of skin cancer^[178–180]. While allicin may have some benefits, there are also potential drawbacks when it comes to using it against skin cancer. Allicin can be a skin irritant and may cause redness, itching, or burning when applied topically. This can be especially problematic for people with sensitive skin or those who are already experiencing skin irritation due to cancer treatments^[182]. Allicin is not widely available in a standardized form for medical use, which can make it difficult to obtain and ensure quality^[183]. It can be concluded that while allicin may have potential benefits for skin cancer, there are also potential drawbacks and limitations to its use.

Sulforaphane

Sulforaphane is a naturally occurring compound found in cruciferous vegetables, such as broccoli, cauliflower, Brussels sprouts, and kale^[184]. Sulforaphane belongs to the family of isothiocyanates. It is formed when the enzyme myrosinase comes into contact with glucoraphanin, a glucosinolate compound present in these vegetables^[185]. Broccoli is the richest dietary source of sulforaphane, with broccoli sprouts containing even higher concentrations of the compound^[186]. Sulforaphane is also available as a dietary supplement, which is typically derived from broccoli sprouts or broccoli seed extract^[187]. It is important to note that the content of sulforaphane in cruciferous vegetables can vary widely depending on factors such as plant variety, growing conditions, and preparation methods. For example, chopping, chewing, or blending cruciferous vegetables can activate myrosinase and increase the formation of sulforaphane, whereas cooking or boiling can reduce the amount of the compound^[188,189].

Sulforaphane has been shown to exhibit chemopreventive and therapeutic effects against various types of cancer, including skin cancer^[190,191]. Its mechanism of action against skin cancer involves several pathways and cellular processes:

Induction of phase II detoxification enzymes

Sulforaphane activates the nuclear factor erythroid 2-related factor 2–antioxidant response element (Nrf2–ARE) signaling pathway, leading to the induction of phase II detoxification enzymes, such as glutathione S-transferases (GSTs), which are involved in the elimination of carcinogens and other toxic compounds from the body^[192].

Inhibition of inflammation

Sulforaphane can modulate the expression of genes involved in inflammation and immune responses, such as NF- κ B and COX-2, leading to a reduction in pro-inflammatory cytokines and chemokines^[193]. Chronic inflammation is known to contribute to the development and progression of skin cancer.

Induction of apoptosis

Sulforaphane can induce apoptosis (programmed cell death) in cancer cells by activating caspases and other apoptotic pathways^[194].

Inhibition of angiogenesis

Sulforaphane can suppress the formation of new blood vessels (angiogenesis) that supply nutrients to cancer cells, thereby inhibiting their growth and proliferation^[195].

Modulation of epigenetic mechanisms

Sulforaphane can modulate epigenetic mechanisms, such as DNA methylation and histone acetylation, leading to changes in gene expression that can affect various cellular processes, including those involved in the development and progression of cancer^[196–198].

Several studies have been conducted by researchers to investigate the molecular mechanism of action of sulforaphane against skin cancer. For instance, Eom *et al.* investigated the effects of sulforaphane treatment on B16F10 melanoma cells and zebrafish models. The results showed that sulforaphane treatment reduced cell proliferation, increased tyrosinase production, and induced cytoskeletal reorganization, leading to an elongated appearance of melanoma cells. Sulforaphane treatment also regulated the protein expression of microphthalmia-associated transcription factor (MITF), protein kinase C beta 1 (PKC β 1), and tyrosinase. The study further demonstrated that sulforaphane-induced biosynthesis of melanin in melanoma cells occurs through changes in actin, as shown by co-treatment of sulforaphane with cytochalasin D (CD) and jasplakinolide (JAS). The same results were obtained in zebrafish models, where sulforaphane upregulated melanin levels despite the presence of the melanin inhibitor phenylthiourea (PTU)^[199]. A study performed by Balasubramanian *et al.* investigated the impact of sulforaphane (SFN), a potential cancer-preventative agent found in cruciferous vegetables, on the expression and function of PcG proteins, which are known to promote cell survival and suppress gene expression in cancer cells. The study found that SFN treatment resulted in a concentration-dependent reduction of PcG protein expression in skin cancer cells, leading to a decrease in histone H3 trimethylation and an accumulation of cells in the G2/M phase. The treatment also increased apoptosis, as evidenced by enhanced cleavage of caspase and PARP proteins. The results suggest that SFN may

inhibit PcG-dependent pro-survival epigenetic events via proteasome-dependent degradation, thus suppressing cancer progression^[196]. In another study, Dickinson *et al.* investigated the chemopreventive properties of sulforaphane, an isothiocyanate found in cruciferous vegetables, against UVB-induced squamous cell carcinoma in mice. The study found that sulforaphane treatment reduced the multiplicity and tumor burden of squamous cell carcinoma in mice co-treated with the carcinogen and sulforaphane. The study also showed that sulforaphane was able to reduce the activity of the transcription factor activator protein-1 (AP-1) in the skin of transgenic mice after UVB. Chromatin immunoprecipitation analysis revealed that sulforaphane inhibited c-Fos, a constituent of the AP-1 dimer, from binding to the AP-1 DNA-binding site. The study also found that sulforaphane and diamide, both known to react with cysteine amino acids, effectively inhibited AP-1 from binding to its response element. Mutation of critical cysteines in the DNA-binding domain of c-Fos and c-Jun resulted in loss of sensitivity to sulforaphane and diamide. These findings suggest that inhibition of AP-1 activity by sulforaphane may be an important mechanism for the chemoprevention of squamous cell carcinoma^[200].

While sulforaphane may have some benefits, there are also potential drawbacks when it comes to using it against skin cancer. The potency of sulforaphane can vary depending on how it is prepared and stored. This can make it difficult to determine the appropriate dosage and ensure consistent results^[201]. Sulforaphane may interact with certain medications, including blood thinners and medications used to treat HIV. Overall, the multiple mechanisms of action of sulforaphane suggest its potential as a promising chemopreventive and therapeutic agent against skin cancer. However, more research is needed to fully understand its effects, safer dose, drug interaction studies and to develop effective treatment strategies.

Ellagic acid

Ellagic acid is a naturally occurring polyphenolic compound found in several fruits and vegetables, including strawberries, raspberries, blackberries, pomegranates, and walnuts^[202]. It is formed from the hydrolysis of ellagitannins, which are water-soluble compounds present in these foods^[203]. Ellagic acid has been shown to exhibit chemopreventive and therapeutic effects against various types of cancer^[204], including skin cancer. Its mechanism of action against skin cancer involves several pathways and cellular processes:

Antioxidant activity

Ellagic acid is a potent antioxidant that can scavenge free radicals and protect cells from oxidative stress-induced damage, which is implicated in the development of skin cancer^[205].

Inhibition of inflammation

Ellagic acid can modulate the expression of genes involved in inflammation and immune responses, such as NF- κ B and COX-2, leading to a reduction in pro-inflammatory cytokines and chemokines^[206]. Chronic inflammation is known to contribute to the development and progression of skin cancer.

Induction of apoptosis

Ellagic acid can induce apoptosis (programmed cell death) in cancer cells by activating caspases and other apoptotic pathways. This mechanism can help eliminate cancer cells and prevent their proliferation^[207–209].

Inhibition of angiogenesis

Ellagic acid can suppress the formation of new blood vessels (angiogenesis) that supply nutrients to cancer cells, thereby inhibiting their growth and proliferation^[123].

Modulation of cellular signaling pathways

Ellagic acid can modulate various signaling pathways involved in the regulation of cell growth, proliferation, and survival, including the MAPK/ERK, PI3K/AKT, and Wnt/ β -catenin pathways. Dysregulation of these pathways is implicated in the development of skin cancer^[67,104].

Researchers have extensively studied the molecular mechanism of action of ellagic acid against skin cancer. Hseu *et al.* investigated the protective effects of ellagic acid against UVA-induced oxidative stress and apoptosis in human keratinocyte cells. Ellagic acid was found to increase cell viability, suppress ROS generation, prevent DNA damage, and inhibit UVA-induced apoptosis. The antioxidant potential of ellagic acid was linked to the increased expression of HO-1 and SOD, downregulation of Keap1, and the activation of Nrf2. Nrf2 knockdown diminished the protective effects of ellagic acid, indicating its potential use for the treatment of UVA-induced skin damage and skin cancer prevention^[204]. In addition, the effects of ellagic acid, a polyphenolic compound from pomegranate fruit extracts, on melanoma cells were investigated. The results showed that ellagic acid significantly inhibited the proliferation, migration, and invasion of WM115 and A375 melanoma cells. Ellagic acid treatment decreased the expression of p-EGFR and vimentin, while it increased the expression of E-cadherin in both cell lines. EGFR activation abolished the effect of ellagic acid on melanoma cells. Additionally, ellagic acid treatment impaired *in vivo* tumorigenesis of A375 cells, and elevated phosphorylated epidermal growth factor receptor (p-EGFR) expression was an independent detrimental factor for melanoma patients. The study suggests that ellagic acid may be useful for the development of new therapeutic strategies for melanoma via the EGFR signaling pathway by Wang *et al.*^[210]. Moreover, Bia *et al.* investigated the stress-resistant action of ellagic acid in *Caenorhabditis elegans* (*C. elegans*) and found that 50 μ M ellagic acid significantly prolonged the lifespan of *C. elegans* under ultraviolet radiation stress, heat stress, oxidative stress, and *Pseudomonas aeruginosa* infection stress. Ellagic acid was also found to reduce damage caused by ultraviolet radiation by inducing the nucleus translocation of Dauer formation 16 (DAF-16) and activating a series of target genes to resist ultraviolet radiation stress. Ellagic acid increased the expression of superoxide dismutase 3 (SOD3) to clean out harmful reactive oxygen species in *C. elegans* exposed to ultraviolet radiation stress. The results suggest that ellagic acid plays an important role in resisting ultraviolet radiation stress in *C. elegans*, probably in an insulin/insulin-like growth factor-1 (IGF-1) signaling pathway-dependent way, and its effects are dependent on the DAF-16 gene, thus helps in preventing skin cancer

and other diseases including diabetes, arteriosclerosis, neurodegenerative diseases, stroke, and cataracts^[211].

While ellagic acid has shown promising health benefits, there are also some potential drawbacks that should be considered. Ellagic acid is poorly absorbed in the body and has low bioavailability. It is rapidly metabolized and eliminated, limiting its potential therapeutic efficacy. Strategies to improve its bioavailability, such as combining it with other compounds or using nanoformulations, are being explored^[212]. Ellagic acid may interact with certain drugs and supplements, including anticoagulants and antiplatelet agents, and may increase the risk of bleeding or bruising. It may also interact with certain chemotherapy drugs and affect their efficacy^[213]. Ellagic acid may cause allergic reactions in some individuals, particularly those who are allergic to berries or other foods that contain the compound^[214]. There is no standardized dosage for ellagic acid, and the optimal dose may vary depending on the specific health condition being targeted^[215]. Additionally, the amount of ellagic acid in foods and supplements can vary widely, making it difficult to determine the actual dose being consumed^[216]. While ellagic acid has shown promising results in preclinical studies, there is limited clinical evidence to support its use in humans. Overall, the multiple mechanisms of action of ellagic acid suggest its potential as a promising chemopreventive and therapeutic agent against skin cancer. However, more research is needed to fully understand its effects, safer dose, drug interaction studies, allergic reaction studies and to develop effective treatment strategies.

Betulinic acid

Betulinic acid is a naturally occurring triterpenoid compound that is found in the bark of several tree species, including white birch (*Betula pubescens*), which is its primary botanical source^[217]. Betulinic acid can also be found in other plants, such as the Chinese herb *Zizyphus jujuba* Mill var. *spinosa*^[218], and in some fruits and vegetables, such as apples^[219] and strawberries^[220], although in lower concentrations. The compound can be extracted from the bark of the white birch tree^[221] using various methods, including maceration and solvent extraction. Betulinic acid has shown a range of biological activities, including anti-inflammatory, antiviral, and anticancer effects^[222,223].

Betulinic acid has been shown to regulate several signaling pathways involved in the development and progression of skin cancer. These pathways include:

PI3K/Akt/mTOR pathway

Betulinic acid can inhibit the PI3K/Akt/mTOR pathway, which is involved in cell growth and survival. This pathway is commonly activated in cancer cells, including skin cancer cells, and contributes to their proliferation and survival^[224].

MAPK/ERK pathway

Betulinic acid can also modulate the MAPK/ERK pathway, which regulates cell proliferation and differentiation. This pathway is often dysregulated in skin cancer and contributes to its development and progression^[225,226].

Wnt/ β -catenin pathway

Betulinic acid has been shown to inhibit the Wnt/ β -catenin pathway, which plays a key role in cell proliferation and stem cell

self-renewal^[227]. Aberrant activation of this pathway has been implicated in the development of various types of cancer, including skin cancer.

NF- κ B pathway

Betulinic acid can also inhibit the NF- κ B pathway, which is involved in inflammation, cell survival, and proliferation. This pathway is commonly activated in cancer cells and contributes to their survival and resistance to chemotherapy^[68].

Several studies have been conducted to investigate the molecular mechanism of action of betulinic acid against skin cancer. For instance, Wróblewska-Łuczka *et al.* investigated the effects of betulinic acid, alone and in combination with taxanes, on the growth of melanoma cell lines. Betulinic acid had no cytotoxic effect on normal cells but significantly inhibited the growth of melanoma cells *in vitro*, with IC₅₀ values ranging from 2.21 to 15.94 μ M. Co-treatment with betulinic acid and taxanes showed desirable drug interactions, with additive and additive with a tendency to synergy interactions observed. These findings suggest that betulinic acid may be a potential therapeutic agent for melanoma, either alone or in combination with taxanes^[228]. One of the studies performed by Liao *et al.* examined the biological effects of betulinic acid (BA)-functionalized GNP in human keratinocytes and melanoma cells. Betulinic acid was grafted onto citrate-capped GNP (BA-GNP) using cysteamine as a linker. The results showed that the BA-GNP formulation had selective cytotoxic and antiproliferative effects on melanoma cells compared to free betulinic acid. Further analysis revealed a pro-apoptotic effect, as evidenced by morphological changes and western blot data showing downregulation of anti-apoptotic Bcl-2 expression and upregulation of pro-apoptotic Bax. GNP also significantly inhibited mitochondrial respiration, demonstrating its mitochondrial-targeted activity. These findings suggest that BA-functionalized GNP could be a potential therapeutic option for melanoma^[229].

The study performed by Kallimanis *et al.* aimed to identify natural compounds that could inhibit aryl hydrocarbon receptor (AhR) activation by these ligands. The methanolic *Rosmarinus officinalis* L. extracts (ROE) were assayed for their activities as antagonists of AhR ligand binding with guinea pig cytosol. The isolated metabolites (*viz.* carnosic acid, carnosol, 7-O-methyl-epi-rosmanol, 4',7-O-dimethylapigenin, and betulinic acid) were assayed for their agonist and antagonist activity in the presence and absence of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) using the gel retardation assay. All assayed extracts showed almost complete inhibition of AhR activation by TCDD at 100 ppm. The methanol ROE at 10 ppm showed significant inhibition against TCDD, 6-formylindolo[3,2-b]carbazole (FICZ), indirubin (IND), and pyrazolo[1,5-a]pyrimidine (PZ), respectively, in human keratinocytes. Most assayed metabolites exhibited dose-dependent antagonist activity. The results suggest that ROE could be useful for the prevention or treatment of skin diseases mediated by the activation of AhR^[230].

Betulinic acid is generally considered safe and non-toxic at therapeutic doses, and it has shown promising anticancer activity in preclinical studies. However, there are some potential drawbacks associated with its use. Betulinic acid has poor water solubility and low bioavailability, which can limit its effectiveness *in vivo*. Various delivery systems and formulations have been developed to enhance its bioavailability and effectiveness^[231–235].

Although betulinic acid has shown promising activity against skin cancer in preclinical studies, there is limited clinical data on its safety and efficacy in humans. Further clinical studies are needed to determine its potential as a therapeutic agent for skin cancer^[236].

Overall, while betulinic acid has shown promise as a potential therapeutic agent against skin cancer, more research is needed to fully understand its bioavailability, safety, efficacy, and optimal use in clinical settings.

Apigenin

Apigenin is a naturally occurring flavone that can be found in various plants, including parsley, chamomile, celery, thyme, and red pepper^[237]. It is also present in many fruits and vegetables, such as oranges, grapefruit, onions, and broccoli. Apigenin can be extracted from these sources using various methods, including microwave-assisted extraction^[238], ultrasound-assisted extraction^[239], supercritical carbon dioxide extraction, enzyme-assisted extraction^[240], high-speed counter-current chromatography^[241], etc. The mechanism of action of apigenin against skin cancer involves multiple pathways, including:

Induction of apoptosis

Apigenin can induce programmed cell death or apoptosis in cancer cells. It does so by activating caspase enzymes, which are responsible for cleaving and degrading proteins that are essential for cell survival. By inducing apoptosis, apigenin can reduce the number of cancer cells in the skin and prevent the formation of tumors^[242].

Inhibition of cell proliferation

Apigenin can inhibit the proliferation of cancer cells by regulating cell cycle progression. It does so by suppressing the expression of cyclin-dependent kinases and upregulating the expression of cell cycle inhibitors. This results in the inhibition of cell division and reduced growth of skin cancer cells^[243].

Regulation of signaling pathways

Apigenin can regulate several signaling pathways that are involved in the development and progression of skin cancer. It can inhibit the activation of the PI3K/Akt^[75,243], MAPK^[112,243], and Wnt/ β -catenin pathways^[244], which promote cell survival, proliferation, and migration. It can also activate the Nrf2/ARE pathway, which is involved in cellular antioxidant and detoxification responses^[245]. By regulating these pathways, apigenin can prevent the growth and spread of skin cancer cells.

Anti-inflammatory effects

Apigenin has been shown to have anti-inflammatory effects that can protect the skin from UV radiation-induced damage. It can reduce the production of pro-inflammatory cytokines and inhibit the activity of enzymes that promote inflammation. By reducing inflammation, apigenin can prevent the formation of skin cancer cells^[246].

Several studies have investigated the molecular mechanism of action of apigenin against skin cancer. In one study conducted by Bridgeman *et al.*, it was found that apigenin inhibited UVB-induced mTOR activation, cell proliferation, and cell cycle

progression in mouse skin and keratinocytes. The inhibition of UVB-induced mTOR signaling by apigenin was not Akt-dependent but instead was driven by adenosine monophosphate-activated protein kinase (AMPK) activation. Additionally, mTOR inhibition by apigenin enhanced autophagy and decreased proliferation in keratinocytes, providing a new target and strategy for better prevention of UV-induced skin cancer^[247]. Another study by Das *et al.* aimed to evaluate the antiproliferative effects of apigenin-loaded poly (lactic-co-glycolide) nanoparticles (NAP) on A375 skin cancer cells *in vitro*. NAP was characterized for particle size, morphology, zeta potential, drug release, and encapsulation. The cellular entry and intracellular localization of NAP were evaluated, along with the stability of dsDNA and relevant markers of mitochondrial functioning, such as ATPase activity, cytochrome-*c* release, and caspase-3 activity. NAP showed better efficacy due to their smaller size and faster mobility with site-specific action. The study revealed that NAP could intercalate with dsDNA, leading to ROS accumulation and depletion of antioxidant enzyme activities, resulting in DNA damage and apoptosis through mitochondrial dysfunction. The study suggests that NAP could be a potential therapeutic option for combating skin melanoma^[248]. A study was performed by Jangdey *et al.* to optimize transfersomes, which are vesicular carriers for drug delivery, using a modified rotary evaporation sonication technique and surfactant Tween 80. The Box-Behnken design with three factors and three levels was applied using response surface methodology. The formulations were characterized for size, shape, entrapment efficiency, stability, and in-vitro permeation. The optimized formulation had an entrapment efficiency of 84.24%, a vesicle size of 35.41 nm, and a drug loading of 8.042%, with good stability. This approach shows promise for the sustained release of apigenin for an extended period of time^[249]. Waheed *et al.* performed a study to develop and optimize lyotropic liquid crystalline nanoparticles (LLC NPs) loaded with apigenin (API) for effective dermal delivery using a quality-by-design (QbD) approach. The optimized API-LLC NPs showed particle size, polydispersity index (PDI), and entrapment efficiency of 287.7 ± 9.53 nm, 0.152 ± 0.051 and 80 ± 2.2 %, respectively. In-vitro and ex-vivo studies showed sustained release and a better permeation profile. The developed API-LLC NPs exhibited better penetration of deeper skin layers, with cytotoxic efficacy assessed on B16F10 cell lines showing a dose-dependent efficacy of API-LLC NPs with an IC_{50} of 45.74 ± 0.05 , making it a promising topical drug delivery nanocarrier for the treatment and management of skin cancer^[250].

Overall, the mechanism of action of apigenin against skin cancer is multifaceted and involves the modulation of several cellular processes. Apigenin is generally considered safe and well-tolerated, and side effects are rare. However, high doses of apigenin supplements or extracts may cause some adverse effects. High doses of apigenin may cause digestive issues such as diarrhea, nausea, and stomach upset^[251]. In some individuals, apigenin may cause an allergic reaction, especially if they have an allergy to other flavonoids or plants in the same family as apigenin^[252]. Apigenin may interact with certain medications, including blood thinners, chemotherapy drugs, and medications that are metabolized by the liver^[253,254]. Although preclinical studies have shown promising results, there is currently a lack of clinical evidence to support the efficacy and safety of apigenin as a treatment for skin cancer. More research is needed to determine its potential benefits and risks.

Gingerol

Gingerol is a bioactive compound found in ginger (*Zingiber officinale*), a spice and medicinal plant that has been used for centuries for its health benefits^[255]. Gingerol belongs to the class of secondary metabolites known as phenolic compounds^[256]. Gingerol is a member of the gingerols, a group of compounds that are responsible for the pungent flavor and aroma of ginger^[257]. Gingerol is known for its anti-inflammatory, antioxidant, and anticancer properties^[258–260]. It has been studied for its potential to help manage a variety of health conditions, including nausea, vomiting, pain, and inflammation^[261,262].

Gingerol has been shown to regulate several signaling pathways involved in the development and progression of skin cancer, including the MAPK/ERK^[83] and PI3K/Akt^[80] pathways. The MAPK/ERK pathway is a signaling pathway that regulates cell growth, division, and survival. Dysregulation of this pathway can contribute to the development of skin cancer. Gingerol has been shown to inhibit the activation of the MAPK/ERK pathway in skin cancer cells, which can help prevent the growth and proliferation of cancer cells^[83]. The PI3K/Akt pathway is another signaling pathway that plays an important role in regulating cell growth, division, and survival. Dysregulation of this pathway has also been implicated in the development of skin cancer. Gingerol has been shown to inhibit the activation of the PI3K/Akt pathway in cancer cells, which can help prevent the growth and proliferation of cancer cells^[80]. In addition to regulating these signaling pathways, gingerol has also been shown to modulate the expression of several genes involved in skin cancer development and progression. For example, gingerol has been shown to upregulate the expression of tumor suppressor genes such as p53 and phosphatase and tensin (PTEN) homolog, which can help prevent the development of skin cancer^[263].

Several studies have been conducted to investigate the molecular mechanism of action of gingerol against skin cancer. A study performed by Praveena *et al.* aimed to investigate the anticancer activity of [6]-gingerol, a bioactive compound found in the rhizome of *Zingiber officinale* and its structural analogs against skin cancer. The ethanolic crude extract of the plant was subjected to phytochemical and gas chromatography–mass spectrometry (GC–MS) analysis to confirm the presence of [6]-gingerol. The anticancer activity was evaluated using the A431 human skin adenocarcinoma cell line, and the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay showed promising cytotoxicity with an IC₅₀ of 81.46 µg/ml. In silico studies were conducted using [6]-gingerol and 21 structural analogs to investigate their anticancer potential and drug-likeness properties. The study targeted the skin cancer protein, DEAD-box helicase 3 X-linked (DDX3X), which regulates all stages of ribonucleic acid (RNA) metabolism, and the compounds were docked to identify the most potent lead molecule based on the lowest binding energy value. The study suggests that [6]-gingerol and its structural analogs could be used as lead molecules for future drug development against skin cancer^[264]. One of the studies conducted by Nigam *et al.* investigated the chemopreventive potential of [6]-gingerol, a pungent ingredient found in ginger rhizome, against benzo[a]pyrene (B[a]P)-induced mouse skin tumorigenesis. Topical treatment of [6]-gingerol was given to animals prior to and after B[a]P treatment. Results showed a delay in tumorigenesis onset, reduced tumor numbers and volume, and elevated apoptotic propensity in tumor tissues.

Western blot analysis showed [6]-gingerol treatment increased p53 levels, Bcl-2-associated X protein (Bax), and apoptotic protease-activating factor-1 (Apaf-1) while decreasing B-cell lymphoma 2 (Bcl-2) and Survivin expression. The study suggests that [6]-gingerol has apoptotic potential as a mechanism of chemoprevention and warrants further investigation^[265]. Another study by Park *et al.* reported the potential cancer chemopreventive properties of [6]-gingerol, a phenolic compound found in ginger, using a two-stage mouse skin carcinogenesis model. The results showed that topical application of [6]-gingerol prior to each dose of 12-O-tetradecanoylphorbol-13-acetate (TPA) significantly inhibited the development of skin papillomagenesis induced by 7,12-dimethylbenz[a]anthracene. The compound also suppressed TPA-induced epidermal inflammation and ornithine decarboxylase activity. These findings suggest that [6]-gingerol may have potential as an antitumor promotional agent^[266].

Gingerol has been associated with several potential health benefits. However, there are also some drawbacks and potential side effects of gingerol. Gingerol may have blood-thinning effects, which can increase the risk of bleeding or interfere with the effectiveness of blood-thinning medications^[267]. Gingerol may interact with certain medications, including anticoagulants, antiplatelets^[267], and blood pressure medications^[268]. Gingerol may not be suitable for pregnant women^[269]. Overall, gingerol appears to regulate multiple signaling pathways involved in the development and progression of skin cancer, which may contribute to its anticancer effects^[270]. However, more research is needed to fully understand the drug interaction studies and safer doses for use in humans.

Quercetin

Quercetin is a natural flavonoid compound found in many fruits, vegetables, and grains^[271]. Some of the food sources of quercetin include onions (*Allium cepa*), apples (*Malus* spp.), berries such as blueberries (*Vaccinium* spp.) and cranberries (*Vaccinium macrocarpon*), citrus fruits such as grapefruit (*Citrus paradisi*), oranges (*Citrus sinensis*), leafy green vegetables such as kale (*Brassica oleracea*) and spinach (*Spinacia oleracea*), and grains such as buckwheat (*Fagopyrum esculentum*)^[272–277]. Flavonoids are responsible for giving plants their vibrant colors^[47]. Quercetin is known for its antioxidant and anti-inflammatory properties and has been studied for its potential health benefits^[278]. The molecular mechanism of action of quercetin against skin cancer involves several pathways and cellular processes. Here are some of the molecular mechanisms by which quercetin acts against skin cancer:

Inhibiting cell proliferation

Quercetin can inhibit cell proliferation by inducing cell cycle arrest at the G1/S phase, which prevents cancer cells from dividing and growing^[58].

Inducing apoptosis

Quercetin can induce apoptosis, or programmed cell death, in cancer cells. This can help eliminate cancer cells and prevent the spread of cancer^[279].

Inhibiting angiogenesis

Quercetin can inhibit the formation of new blood vessels, which is necessary for cancer cells to grow and spread^[87].

Reducing oxidative stress

Quercetin has antioxidant properties and can reduce oxidative stress and damage to skin cells, which can lead to the development of skin cancer^[84].

Regulating signaling pathways

Quercetin can regulate several signaling pathways involved in skin cancer development and progression, including the PI3K/Akt^[279], MAPK^[280], Wnt/ β -catenin^[281], NF- κ B^[282] and JAK/STAT pathways^[270].

Enhancing the immune system

Quercetin can enhance the immune system's ability to recognize and destroy cancer cells, reducing the risk of skin cancer^[84].

In recent years, several studies have investigated the molecular mechanism of action of quercetin against skin cancer. Imran *et al.* developed a nanostructured lipid carrier (NLC) gel loaded with two drugs, quercetin and resveratrol, to enhance their delivery to the dermal and epidermal layers of the skin. The NLC formulation was optimized using a central composite rotatable design (CCRD) and contained a lipid binary mixture, Cremophor RH40 as a surfactant, and had good particle size, polydispersity index, zeta potential, and entrapment efficiency. Dermatokinetic studies showed that the NLC gel significantly increased the disposition of the drugs in the skin compared to a conventional gel, and this was confirmed by confocal microscopic studies. The cytotoxic effect of the NLC gel was assessed in a human epidermoid carcinoma cell line and found to be lower than that of the conventional gel. These results suggest that the NLC gel could be a promising carrier for the delivery of quercetin and resveratrol into deeper layers of the skin for the treatment of skin cancer^[283].

Another study by Caddeo *et al.*, focuses on developing liposomes for delivering two natural polyphenols, quercetin and resveratrol. The liposomes were found to be small, spherical, and uni/bilamellar in nature. The incorporation of polyphenols did not affect their antioxidant activity. Liposomal delivery of polyphenols showed higher cellular uptake and better scavenging ability of ROS in fibroblasts. The in-vivo study in a mouse model of skin lesions demonstrated that topical administration of liposomes reduced tissue damage, edema, and leukocyte infiltration. The study suggests that liposomal delivery of polyphenols may be a promising approach for treating inflammation/oxidative stress associated with precancerous/cancerous skin lesions^[284].

Paliwal *et al.* used ultrasound to enhance the potency of quercetin as a chemotherapeutic drug for the treatment of prostate and skin cancer. The short application of low-frequency ultrasound selectively induced cytotoxicity in cancer cells, while having minimal effect on normal cell lines. The treatment resulted in a significant reduction of viable skin cancer cell population within 48 h. Ultrasound reduced the LC50 of quercetin for skin cancer cells by almost 80-fold, while showing no effect on LC50 for nonmalignant skin cells. The study suggests that ultrasound can be used as a selective sensitizing agent to enhance the efficacy of bioflavonoids for cancer treatment^[285]. Furthermore, Jung *et al.* focus on identifying the molecular targets of quercetin and

its effect on the inhibition of IGF-1 signaling in skin carcinogenesis. The results show that a quercetin diet remarkably delayed the incidence of skin tumor and reduced tumor multiplicity in a mouse skin carcinogenesis protocol. Moreover, skin hyperplasia was significantly inhibited by quercetin supplementation. Further analysis of the skin papilloma cell line showed that quercetin treatment suppressed IGF-1-induced phosphorylation of insulin-like growth factor 1 receptor (IGF-1R), insulin receptor substrate 1 (IRS-1), Akt and ribosomal protein S6 kinase (S6K) and inhibited IGF-1 stimulated cell proliferation. The study suggests that quercetin has potent anticancer activity through the inhibition of IGF-1 signaling and can be considered as a potential therapeutic agent for cancer treatment^[286].

While quercetin has many potential benefits for preventing and treating skin cancer, there are some drawbacks to its use. Quercetin has low bioavailability, which means that much of the compound may be broken down and excreted before it can exert its therapeutic effects^[287]. This can limit its effectiveness against skin cancer. Quercetin may interact with certain medications, such as blood thinners^[288], chemotherapy drugs^[289], and antibiotics^[290]. This can affect their effectiveness and increase the risk of side effects. While preclinical studies have shown promising results for the use of quercetin against skin cancer, there is limited clinical evidence to support its use in humans^[291].

Overall, the molecular mechanisms of action of quercetin against skin cancer are complex and involve multiple cellular processes and pathways. These mechanisms work together to prevent the growth and spread of cancer cells, making quercetin a promising natural compound for skin cancer prevention and treatment. More research is needed to determine its optimal dosage, effectiveness, and safety for preventing and treating skin cancer.

Kaempferol

Kaempferol is a natural flavonoid compound that is found in a variety of plant-based foods such as fruits, vegetables, and herbs^[91]. It is a yellow crystalline solid that belongs to the flavonol subclass of flavonoids^[292]. Kaempferol can be found in plant species like *Ginkgo Folium*^[293]. The molecular mechanism of kaempferol against skin cancer involves several pathways. One of the primary mechanisms is its ability to induce cell cycle arrest and apoptosis in cancer cells. Kaempferol activates the tumor suppressor protein p53, which leads to cell cycle arrest and apoptosis^[294]. Additionally, kaempferol can inhibit the activity of anti-apoptotic proteins such as Bcl-2, which further promotes cell death in cancer cells^[295]. Kaempferol can suppress the production of inflammatory cytokines and chemokines, which can lead to chronic inflammation that promotes the growth of cancer cells^[296]. It can also reduce the generation of reactive oxygen species and enhance the activity of antioxidant enzymes, which helps to protect cells from oxidative damage that can contribute to the development of cancer^[297]. Moreover, kaempferol has been shown to inhibit the activity of several enzymes that are involved in the progression of skin cancer, including tyrosinase^[298], matrix metalloproteinases^[299], and cyclooxygenase-2^[300]. By inhibiting these enzymes, kaempferol can prevent the proliferation and invasion of cancer cells.

Several studies have been performed to elucidate the molecular mechanism of action of kaempferol against skin cancer. In one of the studies, Yang *et al.* evaluated the anticancer activity of

kaempferol against the human malignant melanoma A375 cell line and its effects on apoptosis, cell cycle, cell migration, and mTOR/PI3K/AKT pathway. The results showed that kaempferol exhibited significant anticancer activity against A375 cells with an IC_{50} of 20 μ M. It reduced colony formation in a dose-dependent manner and initiated apoptosis in human malignant melanoma A375 cells. Additionally, kaempferol triggered G2/M cell cycle arrest, inhibited cell migration, and downregulated mTOR, pm-TOR, PI3K, p-PI3K, and Akt protein levels in A375 cells. The findings suggest that kaempferol exerts potent anticancer effects by targeting multiple pathways in melanoma cells^[301]. Yao *et al.* performed a study that investigates the role of the 90 kDa ribosomal S6 kinase (RSK) and mitogen and stress-activated protein kinase (MSK) in solar ultraviolet (SUV) irradiation-induced skin carcinogenesis. The study shows that phosphorylation of RSK and MSK1 is upregulated in human squamous cell carcinoma (SCC) and SUV-treated mouse skin. The study also examines the potential of kaempferol, a natural flavonoid found in tea, broccoli, grapes, apples, and other plant sources, as a chemopreventive agent against SUV-induced skin carcinogenesis. The study reveals that kaempferol inhibits RSK2 and MSK1 kinase activities and, by doing so, attenuates solar UV-induced phosphorylation of cAMP response element-binding protein (CREB) and histone H3 in mouse skin cells. The study further shows that kaempferol is a potent inhibitor of SUV-induced mouse skin carcinogenesis and acts by targeting RSK2 and MSK1. Overall, the study identifies kaempferol as a safe and novel chemopreventive agent against solar UV-induced skin carcinogenesis^[302]. Furthermore, a study by Lee *et al.* investigates the effects of kaempferol, a flavonoid with anti-inflammatory and anti-oxidative properties, on UVB-induced skin inflammation and photocarcinogenesis. The study shows that kaempferol suppresses UVB-induced COX-2 protein expression in mouse skin epidermal JB6 P+ cells and attenuates the UVB-induced transcriptional activities of COX-2 and AP-1. The study further demonstrates that kaempferol attenuates the UVB-induced phosphorylation of several MAPKs, including ERKs, p38, and c-Jun N-terminal kinases (JNKs), by blocking Src kinase activity. The study also shows that kaempferol competes with adenosine triphosphate (ATP) for direct binding to Src and docks easily into the ATP-binding site of Src. The study suggests that kaempferol is a potent chemopreventive agent against skin cancer through its inhibitory interaction with Src. Overall, the study provides insights into the potential of kaempferol as a chemopreventive agent against UVB-induced skin inflammation and photocarcinogenesis^[303].

Overall, the molecular mechanism of kaempferol against skin cancer involves inducing cell cycle arrest and apoptosis, modulating the signaling pathways involved in inflammation and oxidative stress, and inhibiting the activity of enzymes that promote the growth and invasion of cancer cells^[304].

While kaempferol has shown promise in preventing and treating skin cancer, there are some potential drawbacks to its use. Kaempferol has relatively low bioavailability^[305], which means that the body may not absorb it efficiently. Like many natural compounds, kaempferol can have toxic effects at high doses. While it is generally considered safe, some studies have suggested that it can be toxic to certain cells and tissues at high concentrations^[306]. Kaempferol can interact with some medications, particularly those that are metabolized by the liver^[307]. While there have been some promising studies on the use of kaempferol in skin cancer, more research is needed to determine

its safety and effectiveness. It is important to note that most of the research on kaempferol has been done in cell cultures and animal models, and more clinical studies are needed to confirm its effects in humans.

Resveratrol

Resveratrol is a naturally occurring polyphenolic compound found in many plant-based foods, including grapes and red wine^[308]. The botanical name for the grapevine is *Vitis vinifera*, and the most commonly cultivated variety of grapes used for wine production is *Vitis vinifera* subsp. *vinifera*^[309]. Resveratrol is found in the skin of grapes, and its concentration is highest in red grapes compared to white grapes^[310]. Resveratrol is also found in other plants, including peanuts^[311], berries^[312], and knotweed^[313]. The botanical name for the peanut plant is *Arachis hypogaea*, while the most commonly consumed berries that contain resveratrol are blueberries (*Vaccinium* spp.), cranberries (*Vaccinium macrocarpon*), and bilberries (*Vaccinium myrtillus*)^[314]. The Japanese knotweed (*Fallopia japonica*), a plant native to Asia, is also a rich source of resveratrol^[313]. It is classified as a phytoalexin, which means it is produced by plants in response to stress, injury, or infection^[315]. Resveratrol has been extensively studied for its potential health benefits, including its antioxidant^[316], anti-inflammatory^[317], and anticancer^[318] properties. Resveratrol has been shown to have potential anticancer effects against skin cancer through several molecular mechanisms. Here are some of the key ways in which resveratrol may act against skin cancer:

Inhibition of inflammation

Chronic inflammation is a risk factor for the development of many cancers, including skin cancer. Resveratrol has been shown to inhibit the expression of pro-inflammatory cytokines, such as interleukin-1 β and tumor necrosis factor- α , which may help to reduce inflammation and the risk of cancer development^[319,320].

Induction of apoptosis

Resveratrol has been shown to induce apoptosis, or programmed cell death, in skin cancer cells. This is thought to be due in part to its ability to activate certain signaling pathways, such as the p53 pathway, which can trigger cell death^[321,322].

Inhibition of cell proliferation

Resveratrol can also inhibit the proliferation of skin cancer cells, which may help to slow the growth and spread of cancer cells^[96].

Protection against DNA damage

Resveratrol has been shown to have DNA-protective effects, which may help to prevent mutations and other DNA damage that can lead to cancer^[323].

Regulation of cell cycle

Resveratrol can regulate the cell cycle, which is the process by which cells divide and grow. By modulating the activity of certain proteins involved in cell cycle regulation, resveratrol may help to prevent uncontrolled cell growth that can lead to cancer development^[324].

Several studies have investigated the molecular mechanisms underlying the anticancer effects of resveratrol against skin cancer. One study conducted by Osmond *et al.* evaluated the potential of resveratrol as an adjunct to chemotherapy in melanoma treatment. *In vitro*, resveratrol significantly decreased melanoma cell viability in all lines tested and selectively spared nonmalignant fibroblast cell lines. Treatment of malignant cells with resveratrol and temozolomide (TMZ) enhanced cytotoxicity compared to TMZ alone. However, *in vivo*, there was no significant difference. The study suggests that resveratrol has selective antitumor activity *in vitro*, but barriers to translating these results to *in vivo* models need to be overcome^[325]. Another study by Aziz *et al.* discusses the protective role of resveratrol against skin cancer induced by UVR. Resveratrol modulates various molecular mechanisms involved in cell cycle regulation, ROS production, apoptosis, autophagy, cell proliferation, tumor promotion, and cancer-related gene expression. The evidence suggests that resveratrol has effective chemopreventive activity and could potentially be used as a preventive and therapeutic agent in managing UVR-induced skin carcinogenesis^[326]. In addition, resveratrol has been found to modulate the expression of several genes and proteins involved in cell cycle regulation, such as cyclins, cyclin-dependent kinases (CDKs), and checkpoint proteins^[327]. Kundu *et al.* investigate the effects of resveratrol, a compound found in grapes, on COX-2 expression induced by the tumor promoter (TPA) in mouse skin. Results showed that resveratrol significantly inhibited COX-2 expression by suppressing activation of NF- κ B, ERK, and p38 mitogen-activated protein (MAP) kinase. Resveratrol blunted TPA-induced phosphorylation of p65 and its interaction with CBP/p300, rendering NF- κ B transcriptionally inactive. The study also found that resveratrol targets I κ B kinase (IKK) in blocking TPA-induced NF- κ B activation and COX-2 expression in mouse skin *in vivo*^[328]. Furthermore, a study performed by Reagan-Shaw *et al.* investigated the role of cell cycle regulatory molecules in resveratrol-mediated protection against multiple exposures to UVB radiation in hairless mice skin. Resveratrol was topically applied before each UVB exposure, resulting in a decrease in skin thickness, hyperplasia, and leukocyte infiltration. Resveratrol downregulated the upregulation of critical cell cycle regulatory proteins induced by UVB, including proliferating cell nuclear antigen (PCNA), cyclin-dependent kinases, and cyclins. Resveratrol also upregulated cyclin kinase inhibitor WAF1/p21 and tumor suppressor p53, and decreased the upregulation of MAPK, suggesting that the antiproliferative effects of resveratrol may be mediated via modulation of the expression and function of cell cycle regulatory proteins and inhibition of the MAPK pathway. Resveratrol may be useful for preventing UVB-mediated cutaneous damage, including skin cancer^[329]. A study performed by Wu *et al.*, aimed to investigate the effects of resveratrol, a natural polyphenol, on the proliferation and expression of aquaporin 3 (AQP3) in normal human epidermal keratinocytes (NHEKs), which are involved in hyperplastic skin disorders. The results showed that resveratrol inhibited NHEK proliferation in a concentration-dependent manner, and this was associated with downregulation of AQP3 expression and ERK phosphorylation, as well as upregulation of aryl hydrocarbon receptor nuclear translocator (ARNT) expression. The study suggests that the inhibitory effects of resveratrol on AQP3 expression were mediated by Sirtuin 1 (SIRT1)/ARNT/ERK signaling pathway. This

finding may provide insights into the development of new drugs for hyperplastic skin disorders^[330].

Resveratrol's anticancer effects are likely due to its ability to modulate multiple cellular pathways and processes involved in cancer development and progression. While resveratrol has many potential health benefits, including anticancer effects^[318] against skin cancer, there are also some drawbacks and potential risks associated with its use. Resveratrol can interact with certain medications, such as blood thinners^[331], nonsteroidal anti-inflammatory drugs (NSAIDs)^[332], and some antidepressants^[333]. It may also interfere with the effectiveness of chemotherapy drugs^[334]. Resveratrol can cause side effects, including gastrointestinal symptoms (such as nausea, vomiting, and diarrhea)^[335], headaches^[335] and allergic reactions^[336]. Resveratrol supplements are not regulated by the U.S. Food and Drug Administration (FDA), and there is no standardized dosage or quality control. This can make it difficult to ensure the safety and effectiveness of resveratrol supplements^[337]. Overall, while resveratrol has many potential health benefits, it is important to use caution when using it as a supplement, particularly in high doses or in combination with other medications.

Curcumin

Curcumin is a natural compound found in the root of the turmeric plant (*Curcuma longa*), which is a member of the ginger family^[338]. It is a bright yellow pigment and has been used for thousands of years in traditional medicine systems, such as Ayurveda and traditional Chinese medicine, to treat a wide range of conditions^[339]. Curcumin is a polyphenol and is considered to be the most active constituent in turmeric^[340]. It has antioxidant^[341], anti-inflammatory^[342], and anticancer properties^[343] and has been studied extensively for its potential health benefits. It is also a common ingredient in many foods and supplements, including curry powder and turmeric supplements^[338]. In terms of its molecular mechanism against skin cancer, curcumin has been found to inhibit various signaling pathways involved in cancer development and progression. One of the key pathways that curcumin targets is the nuclear factor-kappa B (NF- κ B) signaling pathway^[101]. NF- κ B is a transcription factor that regulates genes involved in inflammation, cell proliferation, and apoptosis, and its activation is commonly found in various types of cancer, including skin cancer. Curcumin has been shown to inhibit NF- κ B activation by suppressing the activity of IKK, which is the kinase that phosphorylates I κ B α and leads to its degradation and subsequent release of NF- κ B from the cytoplasm to the nucleus. Curcumin has also been shown to inhibit the expression of various NF- κ B-regulated genes involved in cell survival, proliferation, and inflammation^[344]. Additionally, curcumin has been found to induce apoptosis (programmed cell death) in skin cancer cells by activating the caspase pathway and inhibiting the anti-apoptotic protein Bcl-2^[345]. Curcumin has also been shown to inhibit angiogenesis (the formation of new blood vessels), which is critical for tumor growth and metastasis^[105].

Multiple investigations have been conducted to explore the molecular mechanisms responsible for the anticancer properties of curcumin against skin cancer. In one of the studies, Zhao *et al.* investigated the effects of curcumin on melanoma cells and found that it effectively inhibited cell proliferation and invasion potential and induced autophagy. Curcumin also arrested the cells at the G2/M phase of the cell cycle and suppressed the activation of AKT, mTOR, and P70S6K proteins, which are part of the AKT/

mTOR signaling pathway. These findings suggest that curcumin could be a potential therapeutic candidate for managing melanoma, which is a highly malignant and resistant cancer^[346]. Another study by Wu *et al.* aimed to investigate the effects of curcumin on the expression of signal transducer and activator of transcription 3 (STAT3) in skin squamous cell carcinoma tissues and the possible mechanisms of curcumin in preventing and treating skin squamous cell carcinoma. Results showed that curcumin treatment inhibited the growth of A431 cells in a time-dependent and dose-dependent manner, with doses above 15 $\mu\text{mol/l}$ for more than 24 h showing significant cytotoxic effects. Curcumin treatment also decreased the invasion and adhesive abilities of A431 cells and inhibited the transcription level of STAT3 mRNA in a dose-dependent manner. The study suggests that curcumin may reduce the invasive ability of A431 cells by inhibiting the activation of the STAT3 signaling pathway and expression of STAT3 as a target gene in the pathway, which may provide insights into the development of new therapies for skin squamous cell carcinoma^[347]. In a study by Kunnumakkara *et al.*, the authors investigated the role of curcumin in the regulation of signaling pathways involved in skin cancer development and progression. The authors demonstrated that curcumin inhibited the expression and activity of various pro-inflammatory cytokines, growth factors, and enzymes involved in skin carcinogenesis, including COX-2, MMP-9, and EGFR. They also showed that curcumin inhibited the activation of the PI3K/Akt/mTOR signaling pathway and the upregulation of its downstream target genes involved in cell survival and proliferation. Furthermore, the authors showed that curcumin enhanced the expression and activity of various tumor suppressor genes, including p53 and p21, which are involved in the regulation of cell cycle progression and apoptosis^[348].

While curcumin has shown promising results in preclinical studies as a potential therapeutic agent against skin cancer, its clinical application has some limitations and challenges. One of the major challenges is its low bioavailability, which refers to the fraction of the administered dose that reaches the systemic circulation in an unchanged form and is available to exert its pharmacological effects. Curcumin is rapidly metabolized in the liver and intestines and has poor solubility in water, which leads to limited absorption and rapid elimination from the body^[349]. Furthermore, curcumin's potential interactions with other drugs and its effects on drug metabolism enzymes have also raised concerns about its safety and efficacy. For instance, curcumin has been reported to inhibit the activity of cytochrome P450 (CYP) enzymes, which play a crucial role in the metabolism of many drugs. This inhibition may result in altered drug efficacy or toxicity when administered in combination with other medications^[350].

Overall, the molecular mechanisms of curcumin against skin cancer are complex and involve multiple pathways and targets. Its ability to modulate inflammation, apoptosis, and angiogenesis makes it a promising natural compound for the prevention and treatment of skin cancer. However, more research is needed to fully understand its bioavailability studies, safer dose, and to optimize its use in clinical settings.

Epigallocatechin gallate

Epigallocatechin gallate is a type of flavonoid, a class of plant-derived compounds that are known for their antioxidant^[351] and

anti-inflammatory properties^[352]. Epigallocatechin gallate is found primarily in green tea, which is derived from the leaves of the *Camellia sinensis* plant^[353]. However, it can also be found in smaller amounts in black tea^[354], white tea^[355], and oolong tea^[356]. Epigallocatechin gallate has been studied for its potential health benefits, including its anticancer properties^[357]. In addition to its anticancer properties, Epigallocatechin gallate has also been studied for its potential role in reducing the risk of cardiovascular disease^[358], improving cognitive function^[359], and promoting weight loss^[358]. It has also been found to have anti-inflammatory effects, which may make it useful in the prevention and treatment of chronic inflammatory diseases^[360] and inflammatory bowel disease^[361]. Epigallocatechin gallate has been shown to have potential chemopreventive and therapeutic effects against skin cancer through several molecular mechanisms of action. These mechanisms include:

Antioxidant activity

Epigallocatechin gallate has strong antioxidant activity, which can protect skin cells from oxidative stress-induced DNA damage, a key contributor to skin cancer development^[106,107].

Anti-inflammatory activity

Epigallocatechin gallate has been found to inhibit the production of pro-inflammatory cytokines and chemokines, which can promote tumor growth and progression. In particular, Epigallocatechin gallate has been shown to inhibit the NF- κ B signaling pathway, which is involved in inflammation and cell survival^[362].

Inhibition of cell proliferation

Epigallocatechin gallate has been found to inhibit the proliferation of skin cancer cells by blocking the cell cycle and inducing apoptosis (programmed cell death). Epigallocatechin gallate has also been found to inhibit the expression of genes involved in cell cycle regulation and DNA repair^[110,111].

Inhibition of angiogenesis

Epigallocatechin gallate has been shown to inhibit angiogenesis, the process by which new blood vessels are formed, which is critical for tumor growth and metastasis^[149,150].

Inhibition of MMPs

Epigallocatechin gallate has been found to inhibit the activity of MMPs, which are enzymes that degrade the extracellular matrix and promote tumor invasion and metastasis^[363].

Numerous studies have been carried out to investigate the molecular mechanisms underlying the anticancer effects of Epigallocatechin gallate in the context of skin cancer. In one of the studies, El-Kayal *et al.* investigated the potential use of ultra-deformable colloidal vesicular systems, including penetration enhancer-containing vesicles, ethosomes, and transethosomes, for the topical delivery of (–)-epigallocatechin-3-gallate as a nutraceutical for skin cancer treatment. The prepared vesicles showed good physical stability and preservation of the antioxidant properties of (–)-epigallocatechin-3-gallate. (–)-epigallocatechin-3-gallate-loaded PEVs and TEs demonstrated an inhibitory effect on epidermoid carcinoma cells and reduced

tumor sizes in mice, with decreased skin oxidative stress biomarkers and lipid peroxidation. These findings suggest that (-)-epigallocatechin-3-gallate PEVs could be an effective topical delivery system for skin cancer treatment^[364]. Zhang *et al.* investigated the interaction between Epigallocatechin-3-gallate and TRAF6 and its effects on the E3 ubiquitin ligase activity of TRAF6, as well as its role in the NF- κ B pathway and melanoma cell growth, migration, and invasion. The results demonstrated that Epigallocatechin-3-gallate binds to TRAF6 and inhibits its E3 ubiquitin ligase activity, leading to the inactivation of the NF- κ B pathway and suppression of melanoma cell growth, migration, and invasion. These findings suggest that Epigallocatechin-3-gallate may be a promising agent for the prevention and treatment of melanoma by targeting TRAF6^[365]. Moreover, Ellis *et al.* describe a study that investigated the mechanism of action of Epigallocatechin-3-gallate, a polyphenolic component of green tea, in inhibiting melanoma cell growth. The study found that Epigallocatechin-3-gallate inhibited melanoma cell growth at physiological doses and that this inhibition was associated with reduced activity of the NF- κ B pathway and decreased secretion of the inflammatory cytokine IL-1 β . The study further demonstrated that Epigallocatechin-3-gallate downregulated the inflammatory component NOD-like receptor (NLR) family pyrin domain-containing 1 (NLRP1). NLRP1 and reduced caspase-1 activation, and silencing NLRP1 expression abolished Epigallocatechin-3-gallate induced inhibition of tumor cell proliferation both *in vitro* and *in vivo*. These findings suggest that inflammasomes and Interleukin-1 beta (IL-1 β) could be potential targets for future melanoma therapeutics^[366].

While Epigallocatechin gallate has shown promise as a potential treatment for skin cancer, there are some drawbacks and limitations to its use. The optimal dose of Epigallocatechin gallate for skin cancer treatment is unclear, and the effective dose may vary depending on the type and stage of the cancer. High doses of Epigallocatechin gallate may also cause adverse effects, such as liver toxicity^[367]. The content of Epigallocatechin gallate in green tea and other natural sources can vary widely, depending on factors such as the plant species, growing conditions, and processing methods. This makes it difficult to standardize the dose and quality of Epigallocatechin gallate in natural sources^[368]. Epigallocatechin gallate has low bioavailability, meaning that it is not well absorbed by the body and has a short half-life. This can limit its effectiveness *in vivo* and make it difficult to achieve therapeutic concentrations in target tissues^[369].

Overall, Epigallocatechin gallate's molecular mechanisms of action against skin cancer involve multiple pathways, including antioxidant^[351], anti-inflammatory^[352], antiproliferative^[370], antiangiogenic^[371], and antimetastatic effects^[371]. These mechanisms suggest that Epigallocatechin gallate may be a promising natural compound for the prevention and treatment of skin cancer, although more research is needed to fully understand its safer dose and potential clinical applications.

β -Carotene

β -Carotene is a red-orange pigment and a type of carotenoid, which is a group of plant pigments responsible for the bright colors of fruits and vegetables^[372]. It is a precursor of vitamin A, meaning that the body can convert it into vitamin A when needed^[373]. β -Carotene is found in high amounts in fruits and vegetables such as carrots, sweet potatoes, pumpkin, spinach, and

apricots^[374]. β -Carotene has antioxidant properties and is believed to have various health benefits, such as reducing the risk of certain types of cancer^[375] and improving immune function^[376].

β -Carotene is obtained from a variety of plants, like carrots (*Daucus carota*), sweet potatoes (*Ipomoea batatas*), pumpkins (*Cucurbita* spp.), etc.^[377]. The molecular mechanism of action of β -carotene against skin cancer involves its ability to neutralize reactive oxygen species (ROS) and reduce oxidative stress, which is a key factor in the development of cancer. ROS are highly reactive molecules that are produced in cells as a result of various physiological and environmental stressors, including exposure to ultraviolet radiation. When ROS levels exceed the capacity of the cell's antioxidant defense system, they can cause oxidative damage to DNA, lipids, and proteins, which can ultimately lead to cancer. β -Carotene can scavenge ROS and neutralize them, thereby reducing the amount of oxidative stress in the cell^[114,115]. Additionally, β -carotene can stimulate the body's natural antioxidant defense system by upregulating the expression of antioxidant enzymes such as superoxide dismutase and catalase. This further helps to reduce oxidative stress and prevent DNA damage^[378]. Moreover, β -carotene also exhibits anti-inflammatory properties that can help reduce chronic inflammation, which is also a risk factor for the development of skin cancer. Chronic inflammation is associated with the production of ROS and DNA damage, which can ultimately lead to the development of cancer^[379].

β -Carotene's ability to scavenge ROS, stimulate antioxidant enzyme expression, and reduce chronic inflammation makes it a potent antioxidant and a promising candidate for the prevention and treatment of skin cancer^[379].

Several researchers have conducted studies on the effect of β -carotene against skin cancer. One such study was conducted by Greenberg *et al.* aimed to investigate whether β -carotene could prevent the occurrence of new nonmelanoma skin cancer in patients who had a recent history of the condition. The study randomly assigned 1805 patients to receive either 50 mg of β -carotene or placebo daily and conducted annual skin examinations. After 5 years, there was no significant difference in the rate of occurrence of new nonmelanoma skin cancer between the groups. The active treatment also showed no efficacy in patients with the lowest initial plasma β -carotene level or in those who currently smoked. The study concludes that β -carotene does not reduce the occurrence of new skin cancers in patients with a previous nonmelanoma skin cancer over a 5-year period of treatment and observation^[380].

In a similar study, Kune *et al.* investigated the relationship between dietary and serum levels of β -carotene, vitamin A, and the risk of nonmelanocytic skin cancer in males. The study found that a high intake of vegetables, including cruciferous vegetables, β -carotene, and vitamin C-containing foods and fish, was inversely related to the risk of skin cancer. The study also found that cases had lower serum levels of β -carotene and vitamin A than controls, and the incidence of skin cancer was inversely related to serum β -carotene levels. Smoking and alcohol consumption did not show a significant association with the risk of skin cancer. The study concludes that further investigation is required to determine the etiologic relevance of low serum levels of β -carotene and vitamin A and the protective effects of a high intake of vegetables and fish on nonmelanocytic skin cancer^[381].

While β -carotene has potential benefits in preventing skin cancer, there are also some drawbacks associated with its use.

Pro-oxidant effects

High doses of β -carotene may have pro-oxidant effects, meaning that they can increase oxidative stress and damage cells. This is particularly true in smokers, as studies have shown that β -carotene supplements can increase the risk of lung cancer in smokers^[382].

Lack of efficacy in some studies

While some studies have suggested that β -carotene can reduce the risk of skin cancer, other studies have shown no significant effect. More research is needed to determine the optimal dose, duration, and form of β -carotene supplementation for skin cancer prevention^[383].

Interactions with other nutrients

β -carotene can interact with other nutrients, such as vitamin E and selenium, which can affect its efficacy^[384]. It is important to ensure adequate intake of these nutrients when supplementing with β -carotene.

Not a substitute for sun protection

β -carotene supplements should not be seen as a substitute for sun protection. While β -carotene may help reduce the risk of skin cancer, it cannot replace the importance of wearing protective clothing and using sunscreen when exposed to ultraviolet radiation^[385].

It can be concluded that while β -carotene may have potential benefits in preventing skin cancer, more research is needed to understand its interaction with other nutrients, which can affect its efficacy.

Caffeic acid

Caffeic acid is a naturally occurring polyphenolic compound and one of the most common hydroxycinnamic acid derivatives^[386]. Caffeic acid is found in a wide variety of plants, including coffee (*Coffea arabica*, *Coffea robusta*), apples (*Malus domestica*), pears (*Pyrus communis*), grapes (*Vitis vinifera*), blueberries (*Vaccinium* spp.), kiwi fruits (*Actinidia deliciosa*), thyme (*Thymus vulgaris*), basil (*Ocimum basilicum*), rosemary (*Rosmarinus officinalis*), sage (*Salvia officinalis*), etc.^[387]. Caffeic acid has been studied for its antioxidant^[388], anti-inflammatory^[389], and anticancer properties^[390]. Caffeic acid has been shown to have potential chemopreventive effects against skin cancer^[390].

The molecular mechanism of action of caffeic acid against skin cancer involves its ability to modulate multiple cellular pathways involved in cell growth, differentiation, and apoptosis. One of the key mechanisms by which caffeic acid exerts its anticancer effects is through the regulation of signaling pathways that control cell proliferation and survival^[126]. Caffeic acid has been shown to inhibit the activity of various enzymes and transcription factors, such as Akt, ERK, NF- κ B^[121,122], and AP-1, which are involved in cell survival and proliferation. By inhibiting these pathways, caffeic acid can induce cell cycle arrest and promote apoptosis, leading to the death of cancer cells^[391–393]. Caffeic acid has also

been shown to possess potent antioxidant and anti-inflammatory properties, which can help to protect cells against oxidative stress and reduce chronic inflammation. Chronic inflammation is a major risk factor for the development of skin cancer, as it can cause DNA damage and impair the immune system's ability to detect and eliminate cancer cells^[394–396]. Additionally, caffeic acid has been shown to enhance the activity of the body's natural defense mechanisms against cancer, such as the activation of the Nrf2 pathway. This pathway is involved in the regulation of antioxidant enzymes and can help to protect cells against oxidative damage^[397–399].

Studies have investigated the molecular mechanisms of action of caffeic acid against skin cancer. One such study was conducted by Yang *et al.*, which investigated the molecular mechanism underlying the anticancer activity of caffeic acid, a phenolic phytochemical found in coffee. The study found that caffeic acid inhibited the colony formation of human skin cancer cells and the neoplastic transformation of HaCaT cells. Topical application of caffeic acid to mouse skin also significantly suppressed tumor incidence and volume in a solar UV-induced skin carcinogenesis model. The study further demonstrated that caffeic acid directly interacted with and inhibited the activity of ERK1 and 2, two proteins involved in mitogen-activated protein kinase signaling. The co-crystal structure of ERK2 complexed with caffeic acid was also resolved, revealing the amino acid residues at which caffeic acid interacts with ERK2. Finally, the study showed that the knockdown of ERK2 in skin cancer cells reduced their sensitivity to caffeic acid in a xenograft mouse model. These findings suggest that caffeic acid exerts its chemopreventive activity against skin carcinogenesis by targeting ERK1 and 2^[400]. Another study by Kudugunti *et al.* investigated the anti-melanoma effects of phenethyl ester of caffeic acid *in vitro* and *in vivo*. The phenethyl ester of caffeic acid was found to inhibit the growth of five different melanoma cell lines, cause intracellular GSH depletion, increase ROS formation, and induce apoptosis in B16-F0 cells. In an *in vivo* study using a B16-F0 melanoma tumor model in mice, it was found to inhibit tumor growth at doses of 10–30 mg/kg/day with minimal toxicity. However, higher doses of phenethyl ester of caffeic acid were associated with increased plasma ALT levels and lipid peroxidation levels in liver and kidney homogenates. The study suggests that phenethyl ester of caffeic acid has potential as an anti-melanoma agent at lower doses^[401]. Furthermore, a study by Balupillai *et al.* investigated the effect of caffeic acid on acute and chronic UVB-irradiation-induced inflammation and photocarcinogenesis in Swiss albino mice. The results showed that caffeic acid administration before UVB exposure decreased inflammation and oxidative stress, enhanced antioxidant status, and activated peroxisome proliferator-activated receptors (PPAR γ) in the mice's skin. PPAR γ is considered a potential target for photo chemoprevention because it inhibits UVB-mediated inflammatory responses. Moreover, both topical and intraperitoneal caffeic acid treatment before each UVB exposure reduced tumor incidence and multiplicity in the mice's skin. Therefore, caffeic acid offers protection against UVB-induced photocarcinogenesis through the activation of the anti-inflammatory transcription factor PPAR γ in mice^[402]. Another study by Balupillai *et al.* investigated the mechanisms by which caffeic acid prevents UVB-induced photocarcinogenesis in human dermal fibroblasts (HDFa) and mouse skin. The results showed that caffeic acid inhibited the formation of cyclobutane pyrimidine dimers (CPDs), oxidative DNA damage, ROS generation,

and apoptotic cell death in HDFa. Furthermore, CA prevented UVB-induced expression of PI3K and AKT kinases through activation of PTEN and promoted XPC-dependent nucleotide excision repair (NER) proteins such as XPC, XPE, transcription factor IIIH (TFIIH-p44), and excision repair cross-complementation group 1 (ERCC1) in human dermal fibroblasts (HDFa) cells and mouse skin tissue. Caffeic acid directly activated PTEN through hydrogen bonds and hydrophobic interactions. These findings suggest that caffeic acid prevents UVB-induced photodamage through the activation of PTEN expression in human dermal fibroblasts and mouse skin^[403]. Another study by Agilan *et al.* investigated the role of JAK-STAT3 signaling in UVB-induced skin carcinogenesis and the protective effect of caffeic acid against it in mouse skin. Chronic UVB irradiation increased the expression of IL-10 and JAK1, which activated STAT3 and led to the transcription of proliferative and anti-apoptotic markers. Caffeic acid inhibited JAK-STAT3 signaling, induced apoptotic cell death, and upregulated the expression of pro-apoptotic markers. Additionally, chronic UVB exposure decreased the expression of thrombospondin-1 (TSP-1), an anti-angiogenic protein, and pretreatment with caffeic acid prevented this loss in UVB-irradiated mouse skin. Thus, caffeic acid offers protection against UVB-induced photocarcinogenesis by modulating the JAK-STAT3 signaling pathway in the mouse skin^[404].

While caffeic acid has shown promising anticancer properties against skin cancer, there are also some potential drawbacks to its use. Some of the possible drawbacks of caffeic acid include:

Dose-dependent toxicity

At high doses, caffeic acid can be toxic to cells and tissues. This can lead to oxidative stress and inflammation, which can actually promote the growth and spread of cancer cells^[405]. Therefore, it is important to use caffeic acid at appropriate doses and under medical supervision.

Bioavailability

Caffeic acid has relatively low bioavailability, which means that a significant amount of it may be metabolized or excreted before it can exert its anticancer effects. To overcome this limitation, researchers are exploring ways to enhance the bioavailability of caffeic acid, such as by using nanoparticles or liposomes^[406].

Lack of clinical trials

While many preclinical studies have shown that caffeic acid has potential anticancer properties, there is a lack of clinical trials to determine its efficacy and safety in humans. More research is needed to establish the optimal dose, route of administration, and duration of treatment^[405].

Interactions with other drugs

Caffeic acid may interact with other drugs or supplements, which could lead to adverse effects or reduce its efficacy. Therefore, it is important to consult a healthcare provider before using caffeic acid as a complementary or alternative therapy for skin cancer^[408].

Overall, the molecular mechanism of action of caffeic acid against skin cancer involves its ability to modulate multiple cellular pathways involved in cell growth, differentiation, and apoptosis. Its potent antioxidant^[388] and anti-inflammatory

properties^[389], along with its ability to enhance the body's natural defense mechanisms, may be used for the prevention and treatment of skin cancer^[390]. However, further research is needed to fully understand its benefits and risks in the prevention and treatment of skin cancer.

Ferulic acid

Ferulic acid is a type of phenolic acid, which is a group of compounds that are widely distributed in plants^[409]. It is a natural antioxidant and has been found in many different plant sources, including fruits, vegetables, grains, and herbs^[410–412]. Ferulic acid is found in many different plant sources, including rice bran (*Oryza sativa*), wheat (*Triticum aestivum*), barley (*Hordeum vulgare*), oats (*Avena sativa*), corn (*Zea mays*), coffee (*Coffea arabica*), pineapple (*Ananas comosus*), apples (*Malus pumila*), oranges (*Citrus sinensis*), and artichokes (*Cynara scolymus*)^[413–415].

Ferulic acid has been shown to have various molecular mechanisms of action against skin cancer. Some of the key mechanisms are:

Antioxidant properties

Ferulic acid has strong antioxidant properties, which means that it can scavenge free radicals and protect skin cells from oxidative stress. This can help to prevent DNA damage and reduce the risk of skin cancer development^[416].

Anti-inflammatory properties

Chronic inflammation is known to contribute to the development of skin cancer. Ferulic acid has been shown to have anti-inflammatory properties, which can help reduce inflammation and prevent the progression of precancerous cells to cancerous cells^[126].

Inhibition of UV-induced damage

Ferulic acid has been shown to inhibit the DNA damage caused by UV radiation, which is a major risk factor for skin cancer. It can also help to repair damaged DNA and prevent the formation of cancerous cells^[417].

Induction of apoptosis

Ferulic acid has been shown to induce apoptosis (programmed cell death) in skin cancer cells. This can help to prevent the growth and spread of cancer cells^[418].

Inhibition of angiogenesis

Ferulic acid has also been shown to inhibit angiogenesis, which is the process by which new blood vessels form to supply nutrients to tumors. By inhibiting angiogenesis, ferulic acid can help to prevent the growth and spread of skin cancer cells^[419].

A study performed by Alias *et al.* aimed to compare the chemopreventive potential of orally administered and topically applied ferulic acid in 7,12-dimethylbenz[a]anthracene (DMBA)-induced skin carcinogenesis. The status of phase I and phase II detoxication agents, lipid peroxidation byproducts, and antioxidants were assessed to determine the mechanistic pathway of their chemopreventive efficacy. Skin squamous cell carcinoma was induced in mice by applying DMBA twice weekly for 8

weeks. Oral administration of ferulic acid completely prevented the formation of skin tumors, whereas topically applied ferulic acid did not show significant chemopreventive activity. Ferulic acid had a modulating effect on the status of lipid peroxidation, antioxidants, and detoxication agents during DMBA-induced skin carcinogenesis. The study concludes that orally administered ferulic acid has a potent suppressing effect on cell proliferation during DMBA-induced skin carcinogenesis^[420]. One study conducted by Choi *et al.* examined the anticancer activity of ferulic acid on National Institutes of Health 3T3 (NIH3T3) fibroblasts and human skin melanoma cells (SK-MEL-3) by measuring the cytotoxicity of ferulic acid on these cells. Ferulic acid decreased cell viability in a dose-dependent manner after human skin melanoma cells were treated with various concentrations of ferulic acid for 48 h. At a concentration of 120 μ M, ferulic acid significantly decreased cell viability in human skin melanoma cells, while it did not show a significant decrease in cell viability at concentrations of 30–120 μ M in NIH3T3 fibroblasts. These results suggest that ferulic acid has anticancer activity in cancer cells, such as human skin melanoma cells, by significantly decreasing cell viability^[421]. Another study by Murakami *et al.* reports the synthesis of a novel ferulic acid analog called FA15, which was found to suppress phorbol ester-induced Epstein–Barr virus activation and superoxide anion generation *in vitro*. The study also demonstrated that FA15 suppressed inducible nitric oxide synthase and cyclooxygenase-2 protein expressions, inhibited the release of tumor necrosis factor- α , and suppressed I- κ B degradation in RAW264.7, a murine macrophage cell line. In mouse skin, topical application of FA15 attenuated hydrogen peroxide production and edema formation, as well as papilloma development, while FA did not. The study concludes that FA15, derived from natural sources, is a novel chemopreventive agent, both structurally and functionally^[422].

While ferulic acid has many potential benefits for the prevention and treatment of skin cancer, there are also some drawbacks and potential side effects. Ferulic acid can cause skin irritation, especially in individuals with sensitive skin. This can manifest as redness, itching, or burning sensations. Ferulic acid can increase the skin's sensitivity to UV radiation, which can increase the risk of sunburn and skin damage. It is important to use sunscreen and limit sun exposure when using ferulic acid-containing skincare products. Ferulic acid can interact with certain medications, including blood thinners and chemotherapy drugs.

Overall, the molecular mechanisms of action of ferulic acid against skin cancer involve its antioxidant^[416] and anti-inflammatory properties^[126], as well as its ability to inhibit UV-induced damage^[417], induce apoptosis^[418], and inhibit angiogenesis^[419]. These properties make ferulic acid a promising natural agent for the prevention and treatment of skin cancer. While ferulic acid is generally considered safe when used in appropriate amounts, it is important to be aware of the potential side effects and to use caution when using it to prevent or treat skin cancer. It is also important to use high-quality, pure products to minimize the risk of skin irritation and other adverse effects.

Importance of a diet rich in fruits and vegetables for preventing skin cancer

A diet rich in fruits and vegetables has been associated with many health benefits, including the prevention of skin cancer^[423]. The

high nutrient content of fruits and vegetables provides the body with the necessary vitamins, minerals, and antioxidants to maintain healthy skin and prevent damage from the sun's harmful UV rays^[424]. One of the most important benefits of a diet rich in fruits and vegetables is its high antioxidant content. Antioxidants are molecules that neutralize free radicals, which are unstable molecules that damage cells and increase the risk of cancer^[425]. Fruits and vegetables contain antioxidants such as vitamin C, vitamin E, and β -carotene, which can help protect the skin from UV radiation and reduce the risk of skin cancer^[426].

Vitamin C is necessary for the production of collagen, a protein that helps to maintain the elasticity of the skin^[427]. β -carotene, which is found in orange and yellow fruits and vegetables, is converted into vitamin A in the body and is important for skin cell growth and repair^[428]. Other vitamins and minerals found in fruits and vegetables, such as potassium and folate, are also essential for maintaining healthy skin^[429]. A diet rich in fruits and vegetables is also high in fiber, which can reduce inflammation in the body^[430]. Chronic inflammation has been linked to an increased risk of many types of cancer, including skin cancer. By reducing inflammation in the body, a diet rich in fruits and vegetables can help prevent the development of skin cancer^[431]. A diet rich in fruits and vegetables can help with weight management^[432]. Being overweight or obese has been linked to an increased risk of certain types of skin cancer, including melanoma. By maintaining a healthy weight, individuals can reduce their risk of developing skin cancer^[433].

To maximize the benefits, individuals should aim to consume a variety of fruits and vegetables in their diet and limit their exposure to UV radiation by wearing protective clothing, seeking shade during peak hours, and using sunscreen^[434].

Fruits and vegetables are rich sources of dietary phytochemicals that play a crucial role in the regulation of cell signaling pathways, which are important in the prevention of skin cancer^[435]. Cell signaling pathways are complex networks of molecular interactions that govern cellular processes such as growth, differentiation, and survival. Abnormalities in these pathways have been linked to the development of skin cancer^[436,437].

Many dietary phytochemicals found in fruits and vegetables that belong to the class of secondary metabolites, like flavonoids, carotenoids, polyphenols, etc., have been shown to regulate cell signaling pathways and prevent the development of skin cancer^[438]. For example, polyphenols found in blueberries have been shown to inhibit the growth of skin cancer cells by regulating the Akt/mTOR signaling pathway^[439]. Flavonoids found in citrus fruits, such as hesperidin and naringenin, have been shown to inhibit the growth of melanoma cells by regulating the MAPK signaling pathway^[440,441].

Fruits and vegetables also have anti-inflammatory properties that can prevent the development of skin cancer by regulating the immune response^[442]. Inflammation is a key component of the skin's response to UV radiation, and chronic inflammation has been linked to an increased risk of skin cancer^[443]. Bioactive compounds found in fruits and vegetables, such as quercetin found in apples and onions, have been shown to inhibit the production of inflammatory cytokines and prevent inflammation in the skin^[444].

It is important to complement dietary changes with other sun safety measures, such as wearing protective clothing, using sunscreen, and limiting sun exposure during peak hours^[445]. This

section includes the dietary sources of some important dietary phytochemicals effective against skin cancer (Table 2A, B).

Blueberries (*Vaccinium corymbosum*)

Blueberries belong to the class of secondary metabolites known as flavonoids^[576]. Specifically, blueberries contain high levels of a type of flavonoid called anthocyanins, which give them their characteristic deep blue color^[577]. Anthocyanins are known for their antioxidant properties^[578], which may have a number of health benefits, including reducing inflammation^[479], improving cardiovascular health^[480], and protecting against certain types of cancer, including skin cancer^[423]. Another phytochemical present in blueberries that may help to prevent skin cancer is pterostilbene^[581], which has been shown to have antioxidant and anti-inflammatory properties and may help to inhibit the growth of skin cancer cells^[582]. Additionally, blueberries are a good source of vitamin C^[583], which has been found to have protective effects against skin damage caused by UV radiation and may also help to prevent skin cancer^[584]. Various research groups have examined its impact on skin cancer. A recent study performed by Alsadi *et al.* examined the effects of a polyphenol-enriched blueberry preparation (PEBP) and non-fermented blueberry juice (NBj) on the expression of miRNAs and target proteins associated with skin cancer. The study found that PEBP was able to

inhibit the proliferation of skin cancer stem cells, reduce the formation of melanophores, and decrease the expression of the CD133+ stem cell marker. The study also found that the expression of tumor suppressor miR-200s was upregulated, and a protein target of the tumor suppressor miR200b, ZEB1, was significantly modulated. These findings suggest that PEBP has potent anticancer and antimetastatic potentials and may represent a novel chemopreventative agent against skin cancer. Qi *et al.* investigated the effects of blueberries on skin cancer prevention in mice. The researchers found that blueberries inhibited skin tumor growth via the regulation of multiple signaling pathways, including the PI3K/AKT, MAPK, and NF-κB pathways. In one of the studies, Afaq *et al.* investigated the effects of green tea and black tea extracts on skin cancer in mice. The researchers found that both green tea and black tea extracts inhibited skin cancer via the regulation of multiple signaling pathways, including the MAPK and PI3K/AKT pathways. Overall, these studies demonstrate that blueberries protect against skin cancer.

Raspberries (*Rubus idaeus*)

Raspberries contain a class of secondary metabolites known as flavonoids, specifically anthocyanins. These are responsible for the fruit's characteristic red color and are also known for their

Table 2
Dietary sources of some important dietary phytochemicals.

S. no.	Dietary phytochemical	Group of plant secondary metabolite	Dietary sources	Reference
A				
1.	Rosmarinic acid	Phenolic acids	Rosemary, sage, thyme, mint, basil, oregano, broccoli, spinach, lemon balm tea, green tea, and some types of honey, particularly those made from plants such as rosemary or thyme.	[446–455]
2.	Allicin	Organosulfur compounds	Raw garlic, onions, leeks, shallots, and chives.	[456–461]
3.	Sulforaphane	Organosulfur compounds	Broccoli, cauliflower, brussels sprouts, kale, cabbage, bok choy, arugula, radishes, and turnips.	[462–470]
4.	Ellagic acid	Polyphenolic compounds	Strawberries, raspberries, blackberries, pomegranates, cranberries, walnuts, pecans, green tea, and black tea.	[471–478]
5.	Betulinic acid	Pentacyclic triterpenoid compounds	Chaga mushrooms, white asparagus, and wild celery.	[479–482]
6.	Apigenin	Flavonoids	Parsley, chamomile, cilantro, thyme, oregano, peppermint, rosemary, celery, onions, artichokes, broccoli, bell peppers, grapefruit, oranges, cherries, chamomile tea, and peppermint tea.	[254,483–495]
7.	Gingerol	Gingerols	Ginger.	[496,497]
8.	Quercetin	Flavonoids	Apples, berries (such as blueberries, cranberries, and elderberries), cherries, grapes, oranges, pomegranates, onions, shallots, kale, broccoli, red leaf lettuce, tomatoes, green peppers, spinach, quinoa, buckwheat, thyme, green tea, and red wine.	[444,498–516]
B				
9.	Kaempferol	Flavonoids	Kale, spinach, broccoli, brussels sprouts, cabbage, grapes, apples, thyme, green tea, black tea, chickpeas and kidney beans.	[517–528]
10.	Resveratrol	Polyphenolic compounds (stilbenes)	Grapes, red wine, blueberries, raspberries, mulberries, peanuts, pistachios, cocoa, dark chocolate, and tomatoes.	[529–533]
11.	Curcumin	Polyphenolic compounds (curcuminoids)	Turmeric.	[534]
12.	Epigallocatechin gallate	Polyphenolic compounds (catechins)	Black tea, green tea (particularly Japanese green tea varieties such as matcha), white tea, strawberries, raspberries, blueberries, grapes, hazelnuts, and pecans.	[354,535–541]
13.	β-Carotene	Terpenoid compounds (carotenoids)	Carrots, sweet potatoes, butternut squash, acorn squash, pumpkin, spinach, kale, honeydew melon, mangoes, and apricots.	[542–550]
14.	Caffeic acid	Polyphenolic compounds (hydroxycinnamic acids)	Coffee, apples, pears, blueberries, cherries, tomatoes, carrots, thyme, oregano, rosemary, sage, almonds, green tea, sunflower seeds, and sesame seeds.	[486,551–562]
15.	Ferulic acid	Polyphenolic compounds (hydroxycinnamic acids)	White bran, germ portions of wheat, rice, oats, oranges, apples, pineapples, cherries, spinach, broccoli, carrots, sweet potatoes, cinnamon, flaxseeds, and sesame seeds.	[386,563–575]

antioxidant properties^[585]. Some studies have suggested that anthocyanins may have protective effects against skin cancer by reducing oxidative stress and inflammation in the skin^[423]. Raspberries are a rich source of dietary phytochemicals, many of which have been found to have potential anticancer properties^[586]. Specifically, several of the compounds found in raspberries have been studied for their ability to prevent or treat skin cancer^[587]. One such compound is ellagic acid, a polyphenol found in raspberries that has been shown to have anti-inflammatory and antioxidant effects^[214]. Studies have found that ellagic acid can inhibit the growth of skin cancer cells and prevent the formation of skin tumors in animal models^[210]. Another phytochemical found in raspberries with potential anticancer properties is quercetin, a flavonoid^[498] with antioxidant^[499] and anti-inflammatory effects^[122]. Quercetin has been shown to induce cell death in skin cancer cells and inhibit the growth of tumors in animal models^[283].

Numerous research teams have looked at its effect on skin cancer. A study performed by Duncan *et al.* investigates the potential of a black raspberry extract (BRE) in reducing cutaneous UVB-induced inflammation and carcinogenesis. Female hairless mice were exposed to UVB radiation and treated topically with either BRE or vehicle control. Results showed that mice treated with BRE had a significant reduction in tumor number and size, which correlated with a decrease in tumor-infiltrating regulatory T-cells. In addition, topical BRE treatment significantly reduced acute UVB-induced inflammation, as evidenced by decreased edema, p53 protein levels, oxidative DNA damage, and neutrophil activation. These findings suggest that BRE may have clinical efficacy in preventing human skin cancers^[588]. Oberyzyzn describes the increased incidence of skin cancer due to exposure to sunlight and changes in tanning practices. Skin tumors are the most common form of cancer in humans, and current treatments for nonmelanoma skin cancers can have severe side effects and limited clinical efficacy. The study proposes natural compounds derived from functional foods as an alternative strategy for the prevention and treatment of skin cancer. Specifically, the study discusses the potential of black raspberry extracts as a chemopreventive/chemotherapeutic agent against nonmelanoma skin cancers^[589]. Mace *et al.* examine the potential of black raspberry extracts (BRB) and their phytochemical metabolites in modulating immune processes relevant to carcinogenesis and immunotherapy. The study found that BRB extracts and their metabolites inhibited the proliferation and viability of CD4+ and CD8+ T lymphocytes, as well as the expansion and suppressive capacity of myeloid-derived suppressor cells (MDSC). Additionally, pretreatment of immune cells with BRB extracts and metabolites attenuated IL-6-mediated phosphorylation of STAT3 and IL-2-induced STAT5 phosphorylation. The study concludes that BRB extracts and their metabolites contain phytochemicals that affect immune processes relevant to carcinogenesis and immunotherapy and could be a potential source of lead compounds for drug development^[590]. Overall, raspberries exhibit promising natural anticancer properties.

Kiwifruit (*Actinidia deliciosa*)

Kiwifruit contains a variety of dietary phytochemicals, including flavonoids, carotenoids, and phenolic acids, that have been studied for their potential benefits in preventing or treating skin

cancer^[591]. Flavonoids, such as quercetin and kaempferol, are particularly abundant in kiwifruit^[592] and have been shown to have antioxidant, anti-inflammatory, and anticancer properties^[593,594]. Studies have suggested that quercetin and kaempferol may be particularly effective against skin cancer due to their ability to inhibit the growth and proliferation of cancer cells, as well as their ability to protect against DNA damage caused by UV radiation^[283,303]. Additionally, the carotenoids present in kiwifruit, such as β -carotene^[595,596], lutein^[596,597], and zeaxanthin^[597], have been studied for their potential to protect against UV-induced skin damage and reduce the risk of skin cancer^[598,599]. Phenolic acids, such as chlorogenic acid, are also present in kiwifruit and have been shown to have antioxidant and anti-inflammatory effects^[600]. Numerous research groups have examined its impact on skin cancer. One of the studies performed by Kou *et al.* investigated the effects of kiwifruit extract on CRL-11147 melanoma cancer cells and the possible mechanisms behind the results. The study found that kiwifruit extract decreased the percentage of cancer cell colonies and increased apoptosis, indicating antitumor effects. The antiproliferative effect of kiwifruit extract was attributed to the downregulation of Cyclin E and CDK4, while the pro-apoptotic effect was attributed to the upregulation of TRAILR1. Overall, the study suggests that kiwifruit extract may have potential for use in the treatment of melanoma^[601]. Another study carried out by Kou *et al.* investigated the potential of kiwifruit extract as a radiosensitizer for Cell Repository Line (CRL-11147) melanoma cancer cells and the underlying mechanisms. Melanoma is a deadly form of skin cancer that is often resistant to radiation therapy. The study used various assays to evaluate the effects of kiwifruit extract on cell proliferation and apoptosis in combination with radiation therapy. The results showed that the combination of kiwifruit extract and radiation therapy decreased the percentage of colonies, Proliferating Cell Nuclear Antigen (PCNA) staining intensity, and optical density value of cancer cells. The study also found that kiwifruit extract increased relative caspase-3 activity, indicating increased apoptosis of cancer cells. The antitumor effect of kiwifruit extract was correlated with increased expression of the antiproliferative molecule p27 and the pro-apoptotic molecule Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand Receptor 1 (TRAILR1). The study suggests that kiwifruit extract could be used as a potential radiosensitizer for melanoma treatment by upregulating both p27 and TRAILR1 to inhibit proliferation and increase apoptosis, respectively^[602]. Overall, these studies show that kiwifruit can help prevent skin cancer.

Watermelon (*Citrullus lanatus*)

Watermelon is a rich source of several dietary phytochemicals that have been found to be effective against skin cancer^[607]. One such phytochemical is lycopene, a carotenoid that gives watermelon its red color. Lycopene is a potent antioxidant that has been shown to protect against UV-induced skin damage, which can lead to skin cancer^[604]. Studies have also found that lycopene can induce cell death in cancer cells and inhibit their growth^[605]. Another phytochemical found in watermelon is citrulline, an amino acid that has been found to have anti-inflammatory properties. Inflammation is a key factor in the development and progression of many types of cancer, including skin cancer. Citrulline has been shown to reduce inflammation and oxidative stress in the skin, potentially reducing the risk of skin cancer^[606].

Watermelon also contains β -carotene, another carotenoid with antioxidant properties. β -Carotene has been shown to protect against UV-induced skin damage and may help reduce the risk of skin cancer^[607]. Its effect on skin cancer has been studied by a plethora of research teams. Ascenso *et al.* focus on the photoprotective properties of lycopene, one of the most potent antioxidants, and its role in anticarcinogenic action at different levels. However, the photoprotective properties of lycopene remain contradictory as some studies show a positive effect while others show a negative effect in both *in vitro* and *in vivo* models. The study highlights the need for a better understanding of the molecular mechanisms of UV damage on skin cells and the role of carotenoids as effective modulators of apoptosis and cell cycle dynamics. The use of lycopene and other effective phyto-compounds as preventive and/or corrective agents may be a potential approach for reducing UVB-induced damage and developing novel therapeutic strategies for skin disorders^[608]. Another study by Fazekas *et al.* investigated the protective effects of topical lycopene, a carotenoid found in tomatoes and watermelon, against acute ultraviolet B (UVB)-induced photodamage. The results showed that the application of lycopene inhibited UVB-induced ornithine decarboxylase and myeloperoxidase and reduced skin thickness. Lycopene also prevented cleavage of caspase-3, restored normal PCNA staining, and maintained normal cell proliferation, suggesting that it may act as a preventative agent by reducing inflammation, maintaining normal cell proliferation, and preventing DNA damage. Thus, topical lycopene showed protective effects against acute UVB-induced photodamage^[609]. Overall, these studies indicate that watermelon can aid in the prevention of skin cancer.

Mangoes (*Mangifera indica*)

Mangoes contain several dietary phytochemicals that have been found to be effective against skin cancer^[610]. One of the primary phytochemicals found in mangoes is quercetin, which has been shown to possess strong antioxidant properties and inhibit the growth of cancer cells^[82,284]. Another important phytochemical found in mangoes is mangiferin, which has been found to possess anti-inflammatory, antioxidant, and anticancer properties^[611]. Studies have shown that mangiferin can inhibit the proliferation of cancer cells and induce apoptosis, which is programmed cell death^[612]. Additionally, β -carotene and lutein, two other important dietary phytochemicals found in mangoes, have been shown to possess photoprotective properties and protect the skin against UV-induced damage, which is a major risk factor for skin cancer^[613,614]. Norathyriol is a flavonoid compound that has been identified in mangoes. It is a naturally occurring phytochemical that belongs to the class of flavanones. Norathyriol has been reported to have various biological activities, including antioxidant, anti-inflammatory, and anticancer properties^[615]. The combination of these dietary phytochemicals found in mangoes makes it a potentially effective dietary intervention for preventing and treating skin cancer. A multitude of research groups have investigated its impact on skin cancer. Li *et al.* describe a study where norathyriol, a compound derived from plants and known to have anticancer activity, was identified through virtual screening of the Chinese Medicine Library as a potential chemopreventive agent for skin cancer. Norathyriol was found to inhibit extracellular signal-regulated kinase (ERK) 1/2 activity and attenuate UVB-induced phosphorylation in

mitogen-activated protein kinases signaling cascades. It was confirmed to bind specifically with ERK2 through co-crystal structural analysis. Norathyriol was shown to inhibit *in vitro* cell growth and significantly suppress solar UV-induced skin carcinogenesis in mouse skin tumorigenesis assays by inhibiting ERK-dependent activity of transcriptional factors AP-1 and NF- κ B during UV-induced skin carcinogenesis. Overall, norathyriol was identified as a safe and highly effective chemopreventive agent against the development of UV-induced skin cancer^[615]. In one of the studies, Saleem *et al.* investigated the antitumor-promoting effects of lupeol, a triterpene found in common fruit plants like mangoes and medicinal herbs, in a mouse skin tumorigenesis model. Topical application of lupeol inhibited conventional and novel biomarkers of tumor promotion in a time-dependent and dose-dependent manner. Lupeol treatment also inhibited the activation of signaling pathways involved in tumor promotion, resulting in reduced tumor incidence, lower tumor burden, and delayed tumor appearance. The results suggest that lupeol is an effective agent for cancer chemoprevention and should be evaluated in tumor models, including skin carcinogenesis^[616]. Overall, the results of these studies suggest that eating mangoes may help lower the risk of getting skin cancer.

Carrots (*Daucus carota*)

Carrots are a rich source of dietary phytochemicals that have been shown to have potential health benefits, including the prevention of skin cancer^[617]. The most notable phytochemicals present in carrots are carotenoids, which include β -carotene, α -carotene, and lutein. These compounds have been shown to have antioxidant properties that protect skin cells from oxidative stress caused by ultraviolet radiation, the most important factor contributing to skin cancer development^[618]. Carotenoids can also enhance immune function and stimulate repair mechanisms in the skin^[619,620]. In addition, falcarinol, a polyacetylene compound found in carrots, has been found to have chemopreventive properties against skin cancer by inducing apoptosis (cell death) in cancer cells and inhibiting their proliferation^[621]. Studies have shown that consuming a diet rich in carrots and other carotenoid-containing vegetables can reduce the risk of developing skin cancer, particularly squamous cell carcinoma^[628]. Therefore, incorporating carrots into a balanced and healthy diet may be an effective strategy for reducing the risk of skin cancer. Numerous research groups have studied its effect on skin cancer. Shebaby *et al.* performed a study that investigates the effects of wild carrot oil fractions on skin cancer. Four fractions were tested for their cytotoxic effects on human epidermal keratinocytes, with one fraction showing a significant antitumor effect on DMBA/TPA-induced skin carcinogenesis in mice. The study also identified a major compound in the fraction, which caused an accumulation of cells in the sub-G1 apoptotic phase and decreased the population of cells in the S and G2/M phases. Additionally, the fraction caused an upregulation of the expression of pro-apoptotic proteins and a downregulation of the expression of anti-apoptotic proteins. The data suggest that the antitumor activity of the fraction may be mediated through inhibition of the MAPK/ERK and PI3K/AKT pathways. Overall, the study suggests that wild carrot oil fractions, particularly the pentane/diethyl ether fraction, may have the potential as a chemopreventive agent against skin cancer^[623]. Natarajmurthy and Dharmesh determined the contribution of phenolics and β -carotene to the antioxidant,

tyrosinase inhibitory, and antiproliferative properties of carrots and evaluated their potential in preventing skin cancer in mice induced by DMBA-UV. Phenolic fractions of carrots were extracted and quantified using high-performance liquid chromatography (HPLC), and their capacities were determined. The results showed that phenolics in CRFP and β -carotene, rather than those in CRBP, exhibited a higher reduction in tumor formation, tyrosinase, and galectin-3 levels and increased anti-oxidant levels, indicating their potential to prevent skin cancer. Thus, carrots enriched with phenolics and β -carotene may be an efficient natural source in preventing skin cancer, as evidenced by the *in vitro* and *in vivo* studies^[624]. In another study, Zeinab *et al.* investigated the chemopreventive effects of oil extract of *Daucus carota* (DCOE) umbels on 7,12-dimethyl benz(a)anthracene (DMBA)-induced skin cancer in mice. The oil extract was administered to animals via gavage, intraperitoneal, and topical routes for 20 weeks, and tumor appearance, incidence, yield, and volume were compared with those of a non-treated control group. The results showed that DCOE has remarkable antitumor activity against DMBA-induced skin cancer compared with non-treated animals, with the topical route being the most effective^[623]. Overall, these studies suggest that carrot consumption may reduce the risk of developing skin cancer.

Future prospects of dietary phytochemicals to prevent skin cancer

The use of dietary phytochemicals as chemopreventive agents in topical skin cancer formulations has gained significant attention in recent years^[435]. While there is promising evidence for the use of phytochemicals in preventing skin cancer, further studies are needed to meet the challenges of topical skin cancer formulations. These challenges include skin penetration, optimum drug concentration, stability, dosing strategy, and sustained drug release following topical application^[625]. Skin penetration is a crucial factor that affects the efficacy of topical formulations^[625]. Dietary phytochemicals, such as flavonoids and polyphenols, have been shown to have low skin penetration due to their large molecular size and low lipophilicity^[627]. Various strategies have been explored to improve the skin penetration of phytochemicals, including the use of nanoemulsions, liposomes, and penetration enhancers^[628]. Optimum drug concentration is another critical factor that needs to be considered in the development of topical skin cancer formulations containing dietary phytochemicals^[625]. The concentration of phytochemicals needs to be optimized to ensure that they are effective in preventing or treating skin cancer while minimizing toxicity. This concentration may vary depending on the phytochemical, the skin type, and the intended use of the formulation^[629]. Stability is also an important factor that needs to be addressed in the development of topical skin cancer formulations containing dietary phytochemicals. The formulation must remain stable during storage and use to ensure that the phytochemicals remain effective^[435]. Stability can be affected by factors such as pH, temperature, and light exposure. Dosing strategy is another factor that needs to be considered in the development of topical skin cancer formulations containing dietary phytochemicals^[435]. The dosing strategy must be optimized to ensure that the phytochemicals are delivered in the most effective manner. Factors such as the frequency of application, the amount of phytochemicals applied, and the duration of treatment

must be carefully considered^[629]. Finally, sustained drug release following topical application is a crucial factor that needs to be addressed in the development of topical skin cancer formulations containing dietary phytochemicals. The phytochemicals must be released in a controlled manner to ensure that they remain effective for the intended duration of treatment^[630]. Various strategies have been explored to achieve sustained drug release, including the use of polymers, liposomes, and nanoparticles^[631]. Overall, while there is promising evidence for the use of dietary phytochemicals in preventing skin cancer, further studies are needed to meet the challenges of topical skin cancer formulations. Addressing these challenges is critical to the development of effective and safe topical formulations containing dietary phytochemicals for the prevention and treatment of skin cancer. The development of such formulations has the potential to revolutionize the management of skin cancer and improve patient outcomes.

Conclusion

Dietary phytochemicals are biologically active compounds that are naturally present in fruits and vegetables. These compounds have been shown to possess numerous health benefits, including the prevention of skin cancer by protecting cells from DNA damage, inhibiting the production of pro-inflammatory cytokines and chemokines, reducing inflammation and the risk of cancer development, inhibiting the activation of signaling pathways (MAPK pathway, PI3K/Akt pathway, Wnt pathway, Hedgehog pathway, Notch pathway, etc.) involved in cell growth and proliferation, which are dysregulated in skin cancer, and inducing apoptosis in cancer cells through various mechanisms like activation of caspases, regulation of Bcl-2 family proteins, modulation of p53, inhibition of NF- κ B, induction of oxidative stress, etc. Assessing the applicability of the preclinical data to humans may require additional research, such as short-term human studies. In addition, studies on skin cancer prevention involving fruits and vegetables seem to be necessary for high-risk humans, such as those with weakened immune systems. In models of high-risk skin carcinogenesis, preclinical studies may reveal positive effects. In addition, fruits and vegetables can be combined with current treatment methods to enhance the treatment of skin cancer.

Ethical approval

Ethics approval was not required for this review.

Consent

Informed consent was not required for this review.

Source of funding

No funding.

Author contribution

H.S. and A.K.M.: conceptualization, data curation, writing – original draft preparation, and writing – reviewing and editing; S.M., A.K., A.M., and R.A.: data curation, writing – original

draft preparation, and writing – reviewing and editing; C.R.D.: writing – reviewing and editing and visualization; T.B.E.: writing – reviewing and editing, visualization, and supervision.

Conflicts of interest disclosure

All authors report no conflicts of interest relevant to this article.

Research registration unique identifying number (UIN)

1. Name of the registry: not applicable.
2. Unique identifying number or registration ID: not applicable.
3. Hyperlink to your specific registration (must be publicly accessible and will be checked): not applicable.

Guarantor

Talha Bin Emran, PhD, Associate Professor, Department of Pharmacy, BGC Trust University Bangladesh, Chittagong 4381, Bangladesh; Tel.: +88 030 3356193; fax: +88 031 2550224; cell: +88 01819 942214; <https://orcid.org/0000-0003-3188-2272>; e-mail: talhabmb@bgctub.ac.bd.

Data availability statement

Not applicable to this article.

Provenance and peer review

Not commissioned, externally peer-reviewed.

References

- [1] Proksch E, Brandner JM, Jensen JM. The skin: an indispensable barrier. *Exp Dermatol* 2008;17:1063–72.
- [2] Elias PM, Williams ML. Overview of the epidermis and dermis. In: Goldsmith S, Wolff K, Katz S, Gilchrist B, Paller A, Leffell D, eds. *Fitzpatrick's Dermatology In General Medicine*, 9th ed. McGraw-Hill Education; 2019. pp. 61–72.
- [3] Madison KC. Barrier function of the skin: “La raison d’être” of the epidermis. *J Invest Dermatol* 2003;121:231–41.
- [4] Losquadro WD. Anatomy of the skin and the pathogenesis of non-melanoma skin cancer. *Facial Plast Surg Clin North Am* 2017;25:283–9.
- [5] Driskell RR, Watt FM. Understanding fibroblast heterogeneity in the skin. *Trends Cell Biol* 2015;25:92–9.
- [6] Nguyen AV, Soulika AM. The dynamics of the skin's immune system. *Int J Mol Sci* 2019;20:1811.
- [7] Senechal B, Elain G, Jezierski E, *et al.* Expansion of regulatory T cells in patients with Langerhans cell histiocytosis. *PLoS Med* 2007;4:e253.
- [8] Rogers HW, Weinstock MA, Feldman SR, *et al.* Incidence estimate of nonmelanoma skin cancer (keratinocyte carcinomas) in the U.S. population, 2012. *JAMA Dermatol* 2015;151:1081–6.
- [9] Leiter U, Eigentler T, Garbe C. Epidemiology of skin cancer. *Adv Exp Med Biol* 2014;810:120–40.
- [10] Stratigos AJ, Garbe C, Dessinioti C, *et al.* European interdisciplinary guideline on invasive squamous cell carcinoma of the skin: Part 2. Treatment. *Eur J Cancer* 2020;128:83–102.
- [11] Kwong LN, Davies MA. Navigating the therapeutic complexity of PI3K pathway inhibition in melanoma. *Clin Cancer Res* 2013;19:5310–9.
- [12] Lomas A, Leonardi-Bee J, Bath-Hextall F. A systematic review of worldwide incidence of nonmelanoma skin cancer. *Br J Dermatol* 2012;166:1069–80.
- [13] Epstein EH. Basal cell carcinomas: attack of the hedgehog. *Nat Rev Cancer* 2008;8:743–54.
- [14] Wong SY, Reiter JF. The primary cilium at the crossroads of mammalian hedgehog signalling. *Curr Top Dev Biol* 2008;85:225–60.
- [15] Tang JY, Aszterbaum M, Athar M, *et al.* Basal cell carcinoma chemoprevention with nonsteroidal anti-inflammatory drugs in genetically predisposed PTCH1+/- humans and mice. *Cancer Prev Res (Phila)* 2010;3:25–34.
- [16] Yin VT, Esmaili B. Targeting the Hedgehog pathway for locally advanced and metastatic basal cell carcinoma. *Curr Pharm Des* 2017;23:655–9.
- [17] Rivlin N, Brosh R, Oren M, *et al.* Mutations in the p53 tumor suppressor gene: important milestones at the various steps of tumorigenesis. *Genes Cancer* 2011;2:466–74.
- [18] Mohan SV, Chang AL. Advanced basal cell carcinoma: epidemiology and therapeutic innovations. *Curr Dermatol Rep* 2014;3:40–5.
- [19] Yanofsky VR, Mercer SE, Phelps RG. Histopathological variants of cutaneous squamous cell carcinoma: a review. *J Skin Cancer* 2011;2011:210813.
- [20] Olivier M, Hollstein M, Hainaut P. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol* 2010;2:a001008.
- [21] Weissmueller S, Manchado E, Saborowski M, *et al.* Mutant p53 drives pancreatic cancer metastasis through cell-autonomous PDGF receptor beta signalling. *Cell* 2014;157:382–94.
- [22] Harismendy O, Notani D, Song X, *et al.* 9p21 DNA variants associated with coronary artery disease impair interferon-gamma signalling response. *Nature* 2011;470:264–8.
- [23] Stiewe T, Zimmermann S, Frilling A, *et al.* Transactivation-deficient DeltaTA-p73 acts as an oncogene. *Cancer Res* 2002;62:3598–602.
- [24] Basu TN, Gutmann DH, Fletcher JA, *et al.* Aberrant regulation of ras proteins in malignant tumour cells from type 1 neurofibromatosis patients. *Nature* 1992;356:713–5.
- [25] Schubert S, Shannon K, Bollag G. Hyperactive Ras in developmental disorders and cancer. *Nat Rev Cancer* 2007;7:295–308.
- [26] Hodis E, Watson IR, Kryukov GV, *et al.* A landscape of driver mutations in melanoma. *Cell* 2012;150:251–63.
- [27] Dankort D, Curley DP, Carlidge RA, *et al.* Braf(V600E) cooperates with Pten loss to induce metastatic melanoma. *Nat Genet* 2009;41:544–52.
- [28] Chin L, Garraway LA, Fisher DE. Malignant melanoma: genetics and therapeutics in the genomic era. *Genes Dev* 2006;20:2149–82.
- [29] Lin WM, Baker AC, Beroukhim R, *et al.* Modeling genomic diversity and tumor dependency in malignant melanoma. *Cancer Res* 2008;68:664–73.
- [30] Viros A, Fridlyand J, Bauer J, *et al.* Improving melanoma classification by integrating genetic and morphologic features. *PLoS Med* 2008;5:e120.
- [31] Hocker T, Tsao H. Ultraviolet radiation and melanoma: a systematic review and analysis of reported sequence variants. *Hum Mut* 2007;28:578–88.
- [32] Hayward NK, Wilmott JS, Waddell N, *et al.* Whole-genome landscapes of major melanoma subtypes. *Nature* 2017;545:175–80.
- [33] Shain AH, Yeh I, Kovalyshyn I, *et al.* The genetic evolution of melanoma from precursor lesions. *N Engl J Med* 2015;373:1926–36.
- [34] Oren M, Levine AJ. Molecular cloning of a cDNA specific for the murine p53 cellular tumor antigen. *Proc Natl Acad Sci U S A* 1983;80:56–9.
- [35] Soussi T, Wiman KG. Shaping genetic alterations in human cancer: the p53 mutation paradigm. *Cancer Cell* 2007;12:303–12.
- [36] Vogelstein B, Lane D, Levine AJ. Surfing the p53 network. *Nature* 2000;408:307–10.
- [37] Freed-Pastor WA, Prives C. Mutant p53: one name, many proteins. *Genes Dev* 2012;26:1268–86.
- [38] Toledo F, Wahl GM. Regulating the p53 pathway: in vitro hypotheses, in vivo veritas. *Nat Rev Cancer* 2006;6:909–23.
- [39] Li Y, Zhang T, Korkaya H, *et al.* Sulforaphane, a dietary component of broccoli/broccoli sprouts, inhibits breast cancer stem cells. Sulforaphane inhibits breast cancer stem cells. *Clin Cancer Res* 2010;16:2580–90.
- [40] Patil JR, Jayaprakash GK, Murthy KC, *et al.* Apoptosis-mediated proliferation inhibition of human colon cancer cells by volatile principles of *Citrus aurantifolia*. *Food Chem* 2009;114:1351–8.
- [41] Grosso G, Bei R, Mistretta A, *et al.* Effects of vitamin C on health: a review of evidence. *Front Biosci (Landmark Ed)* 2013;18:1017–29.
- [42] Spencer JP, Vafeiadou K, Williams RJ, *et al.* Neuroinflammation: modulation by flavonoids and mechanisms of action. *Mol Aspects Med* 2012;33:83–97.

- [43] Pandey KB, Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid Med Cell Longev* 2009;2:270–8.
- [44] Johnson IT. Phytochemicals and cancer. *Proc Nutr Soc* 2007;66:207–15.
- [45] Manach C, Scalbert A, Morand C, *et al.* Polyphenols: food sources and bioavailability. *Am J Clin Nutr* 2004;79:727–47.
- [46] Liu RH. Health-promoting components of fruits and vegetables in the diet. *Adv Nutr* 2013;4:384S–92S.
- [47] Kesarkar S, Bhandage A, Deshmukh S, *et al.* Flavonoids: an overview. *J Pharm Res* 2009;2:1148–54.
- [48] Grosso G, Godos J, Lamuela-Raventos R, *et al.* A comprehensive meta-analysis on dietary flavonoid and lignan intake and cancer risk: level of evidence and limitations. *Mol Nutr Food Res* 2017;61:1600930.
- [49] Stahl W, Sies H. Antioxidant activity of carotenoids. *Mol Aspects Med* 2003;24:345–51.
- [50] Tanaka T, Shnimizu M, Moriwaki H. Cancer chemoprevention by caroteno. *Molecules* 2012;17:3202–42.
- [51] Coulman KD, Liu Z, Hum WQ, *et al.* Whole sesame seed is as rich a source of mammalian lignan precursors as whole flaxseed. *Nutr Cancer* 2005;52:156–65.
- [52] Adlercreutz H. Lignans and human health. *Crit Rev Clin Lab Sci* 2007;44:483–525.
- [53] Shiau JP, Chuang YT, Tang JY, *et al.* The impact of oxidative stress and AKT pathway on cancer cell functions and its application to natural products. *Antioxidants (Basel)* 2022;11:1845.
- [54] Sana SS, Pasha A, Kumbhakar DV, *et al.* SnO₂-NPs (synthesized from crude rosmarinic acid) inhibit HeLa cells progression in vitro for anti-tumor investigation and act as an angiogenic inhibitor in ovo CAM xenograft. *Nano-Struct Nano-Objects* 2023;33:100918.
- [55] Eskandani R, Kazempour M, Farahzadi R, *et al.* Engineered nanoparticles as emerging gene/drug delivery systems targeting the nuclear factor-κB protein and related signalling pathways in cancer. *Biomed Pharmacother* 2022;156:113932.
- [56] Yuan Y, Guo M, Gu C, *et al.* The role of Wnt/β-catenin signalling pathway in the pathogenesis and treatment of multiple myeloma (review). *Am J Transl Res* 2021;13:9932–49.
- [57] Zhang X, Xie K, Zhou H, *et al.* Role of non-coding RNAs and RNA modifiers in cancer therapy resistance. *Mol Cancer* 2020;19:1–26.
- [58] Gupta SC, Kim JH, Prasad S, *et al.* Regulation of survival, proliferation, invasion, angiogenesis, and metastasis of tumor cells through modulation of inflammatory pathways by nutraceuticals. *Cancer Metastasis Rev* 2010;29:405–34.
- [59] Mondal A, Banerjee S, Bose S, *et al.* Garlic constituents for cancer prevention and therapy: From phytochemistry to novel formulations. *Pharmacol Res* 2022;175:105837.
- [60] Huang Y, Li W, Su ZY, *et al.* The complexity of the Nrf2 pathway: beyond the antioxidant response. *J Nutr Biochem* 2015;26:1401–13.
- [61] Roy SK, Srivastava RK, Shankar S. Inhibition of PI3K/AKT and MAPK/ERK pathways causes activation of FOXO transcription factor, leading to cell cycle arrest and apoptosis in pancreatic cancer. *J Mol Signal* 2010;5:10.
- [62] Wang L, Tian Z, Yang Q, *et al.* Sulforaphane inhibits thyroid cancer cell growth and invasiveness through the reactive oxygen species-dependent pathway. *Oncotarget* 2015;6:25917.
- [63] Ołędzka AJ, Czerwińska ME. Role of plant-derived compounds in the molecular pathways related to inflammation. *Int J Mol Sci* 2023;24:4666.
- [64] Edderkaoui M, Odinkova I, Ohno I, *et al.* Ellagic acid induces apoptosis through inhibition of nuclear factor κB in pancreatic cancer cells. *World J Gastroenterol* 2008;14:3672.
- [65] Zhang HM, Zhao L, Li H, *et al.* Research progress on the anticarcinogenic actions and mechanisms of ellagic acid. *Cancer Biol Med* 2014;11:92.
- [66] Qiu S, Zhong C, Zhao B, *et al.* Transcriptome analysis of signalling pathways targeted by Ellagic acid in hepatocellular carcinoma cells. *Biochim Biophys Acta Gen Subj* 2021;1865:129911.
- [67] Ceci C, Lacal PM, Tentori L, *et al.* Experimental evidence of the anti-tumor, antimetastatic and antiangiogenic activity of ellagic acid. *Nutrients* 2018;10:1756.
- [68] Takada Y, Aggarwal BB. Betulinic acid suppresses carcinogen-induced NF-κB activation through inhibition of IκBα kinase and p65 phosphorylation: abrogation of cyclooxygenase-2 and matrix metalloproteinase-9. *J Immunol* 2003;171:3278–86.
- [69] Mullauer FB, Kessler JH, Medema JP. Betulinic acid, a natural compound with potent anticancer effects. *Anticancer Drugs* 2010;21:215–27.
- [70] Fulda S, Friesen C, Los M, *et al.* Betulinic acid triggers CD95 (APO-1/Fas)-and p53-independent apoptosis via activation of caspases in neuroectodermal tumors. *Cancer Res* 1997;57:4956–64.
- [71] Jiang W, Li X, Dong S, *et al.* Betulinic acid in the treatment of tumour diseases: application and research progress. *Biomed Pharmacother* 2021;142:111990.
- [72] Liu W, Li S, Qu Z, *et al.* Betulinic acid induces autophagy-mediated apoptosis through suppression of the PI3K/AKT/mTOR signalling pathway and inhibits hepatocellular carcinoma. *Am J Transl Res* 2019;11:6952.
- [73] Jiang N, Hu Y, Wang M, *et al.* The Notch signalling pathway contributes to angiogenesis and tumor immunity in breast cancer. *Breast Cancer (Dove Med Press)* 2022;14:291–309.
- [74] Tong X, Pelling JC. Targeting the PI3K/Akt/mTOR axis by apigenin for cancer prevention. *Anticancer Agents Med Chem* 2013;13:971–8.
- [75] Javed Z, Sadia H, Iqbal MJ, *et al.* Apigenin role as cell-signalling pathways modulator: implications in cancer prevention and treatment. *Cancer Cell Int* 2021;21:189.
- [76] Zhao G, Han X, Cheng W, *et al.* Apigenin inhibits proliferation and invasion, and induces apoptosis and cell cycle arrest in human melanoma cells. *Oncol Rep* 2017;37:2277–85.
- [77] Jia M, Mateoiu C, Souchelnytskyi S. Protein tyrosine nitration in the cell cycle. *Biochem Biophys Res Commun* 2011;413:270–6.
- [78] Shukla S, Gupta S. Apigenin: a promising molecule for cancer prevention. *Pharm Res* 2010;27:962–78.
- [79] Imran M, Aslam Gondal T, Atif M, *et al.* Apigenin as an anticancer agent. *Phytother Res* 2020;34:1812–28.
- [80] Deng J, Bai X, Feng X, *et al.* Inhibition of PI3K/Akt/mTOR signalling pathway alleviates ovarian cancer chemoresistance through reversing epithelial–mesenchymal transition and decreasing cancer stem cell marker expression. *BMC Cancer* 2019;19:618.
- [81] Kim SO, Kundu JK, Shin YK, *et al.* [6]-Gingerol inhibits COX-2 expression by blocking the activation of p38 MAP kinase and NF-κB in phorbol ester-stimulated mouse skin. *Oncogene* 2005;24:2558–67.
- [82] Surh YJ, Na HK. NF-κB and Nrf2 as prime molecular targets for chemoprevention and cytoprotection with anti-inflammatory and antioxidant phytochemicals. *Genes Nutr* 2008;2:313–7.
- [83] Huang HC, Chou YC, Wu CY, *et al.* [8]-Gingerol inhibits melanogenesis in murine melanoma cells through down-regulation of the MAPK and PKA signal pathways. *Biochem Biophys Res Commun* 2013;438:375–81.
- [84] Rather RA, Bhagat M. Quercetin as an innovative therapeutic tool for cancer chemoprevention: molecular mechanisms and implications in human health. *Cancer Med* 2020;9:9181–92.
- [85] Xiong G, Ji W, Wang F, *et al.* Quercetin inhibits inflammatory response induced by LPS from *Porphyromonas gingivalis* in human gingival fibroblasts via suppressing NF-κB signalling pathway. *Biomed Res Int* 2019;2019:6282635.
- [86] Zehsaz F, Farhangi N, Mirheidari L. The effect of *Zingiber officinale* R. rhizomes (ginger) on plasma pro-inflammatory cytokine levels in well-trained male endurance runners. *Cent Eur J Immunol* 2014;39:174–80.
- [87] Zalpoor H, Nabi-Afjadi M, Forghanifidvajani R, *et al.* Quercetin as a JAK-STAT inhibitor: a potential role in solid tumors and neurodegenerative diseases. *Cell Mol Biol Lett* 2022;27:60.
- [88] Sekiguchi A, Motegi SI, Fujiwara C, *et al.* Inhibitory effect of kaempferol on skin fibrosis in systemic sclerosis by the suppression of oxidative stress. *J Dermatol Sci* 2019;96:8–17.
- [89] Sharma V, Joseph C, Ghosh S, *et al.* Kaempferol induces apoptosis in glioblastoma cells through oxidative stress. *Mol Cancer Ther* 2007;6:2544–53.
- [90] Ashrafzadeh M, Gholami MH, Mirzaei S, *et al.* Dual relationship between long non-coding RNAs and STAT3 signalling in different cancers: new insight to proliferation and metastasis. *Life Sci* 2021;270:119006.
- [91] Chen AY, Chen YC. A review of the dietary flavonoid, kaempferol on human health and cancer chemoprevention. *Food Chem* 2013;138:2099–107.
- [92] Kim GD. Kaempferol inhibits angiogenesis by suppressing HIF-1α and VEGFR2 activation via ERK/p38 MAPK and PI3K/Akt/mTOR signalling pathways in endothelial cells. *Prevent Nutr Food Sci* 2017;22:320.

- [93] Chikara S, Nagaprashantha LD, Singhal J, *et al.* Oxidative stress and dietary phytochemicals: role in cancer chemoprevention and treatment. *Cancer Lett* 2018;413:122–34.
- [94] Kundu JK, Shin YK, Surh YJ. Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin in vivo: NF- κ B and AP-1 as prime targets. *Biochem Pharmacol* 2006;72:1506–15.
- [95] Nichols JA, Katiyar SK. Skin photoprotection by natural polyphenols: anti-inflammatory, antioxidant and DNA repair mechanisms. *Arch Dermatol Res* 2010;302:71–83.
- [96] Wu Z, Liu B, E C, *et al.* Resveratrol inhibits the proliferation of human melanoma cells by inducing G1/S cell cycle arrest and apoptosis. *Mol Med Rep* 2015;11:400–4.
- [97] Aziz MH, Reagan-Shaw S, Wu J, *et al.* Chemoprevention of skin cancer by grape constituent resveratrol: relevance to human disease? *FASEB J* 2005;19:1193–5.
- [98] Elshaer M, Chen Y, Wang XJ, *et al.* Resveratrol: an overview of its anti-cancer mechanisms. *Life Sci* 2018;207:340–9.
- [99] Nakamura Y, Ohto Y, Murakami A, *et al.* Inhibitory effects of curcumin and tetrahydrocurcuminoids on the tumor promoter-induced reactive oxygen species generation in leukocytes in vitro and in vivo. *Jpn J Cancer Res* 1998;89:361–70.
- [100] He Y, Yue Y, Zheng X, *et al.* Curcumin, inflammation, and chronic diseases: how are they linked? *Molecules* 2015;20:9183–213.
- [101] Chao SC, Hu DN, Roberts J, *et al.* Inhibition effect of curcumin on UVB-induced secretion of pro-inflammatory cytokines from corneal limbus epithelial cells. *Int J Ophthalmol* 2017;10:827.
- [102] Yu D, An F, He X, *et al.* Curcumin inhibits the proliferation and invasion of human osteosarcoma cell line MG-63 by regulating miR-138. *Int J Clin Exp Pathol* 2015;8:14946.
- [103] Arbiser JL, Klaubner N, Rohan R, *et al.* Curcumin is an in vivo inhibitor of angiogenesis. *Mol Med* 1998;4:376–83.
- [104] Panahi Y, Fazlollahzadeh O, Atkin SL, *et al.* Evidence of curcumin and curcumin analogue effects in skin diseases: a narrative review. *J Cell Physiol* 2019;234:1165–78.
- [105] Bhandarkar SS, Arbiser JL. Curcumin as an inhibitor of angiogenesis. *Adv Exp Med Biol* 2007;595:185–95.
- [106] Huang CC, Wu WB, Fang JY, *et al.* (-)-Epicatechin-3-gallate, a green tea polyphenol is a potent agent against UVB-induced damage in HaCaT keratinocytes. *Molecules* 2007;12:1845–58.
- [107] Salaheldin TA, Adhamsi VM, Fujioka K, *et al.* Photochemoprevention of ultraviolet beam radiation-induced DNA damage in keratinocytes by topical delivery of nanoformulated epigallocatechin-3-gallate. *Nanomedicine* 2022;44:102580.
- [108] Kim I, He YY. Ultraviolet radiation-induced non-melanoma skin cancer: regulation of DNA damage repair and inflammation. *Genes Dis* 2014;1:188–98.
- [109] Sharifi-Rad M, Pezzani R, Redaelli M, *et al.* Preclinical activities of epigallocatechin gallate in signalling pathways in cancer. *Molecules* 2020;25:467.
- [110] Aggarwal V, Tuli HS, Tania M, *et al.* Molecular mechanisms of action of epigallocatechin gallate in cancer: recent trends and advancement. *Semin Cancer Biol* 2022;80:256–75.
- [111] Choudhury SR, Balasubramanian S, Chew YC, *et al.* (-)-Epigallocatechin-3-gallate and DZNep reduce polycomb protein level via a proteasome-dependent mechanism in skin cancer cells. *Carcinogenesis* 2011;32:1525–32.
- [112] Katiyar S, Elmets CA, Katiyar SK. Green tea and skin cancer: photo-immunology, angiogenesis and DNA repair. *J Nutr Biochem* 2007;18:287–96.
- [113] Jung YD, Ellis LM. Inhibition of tumour invasion and angiogenesis by epigallocatechin gallate (EGCG), a major component of green tea. *Int J Exp Pathol* 2001;82:309–16.
- [114] O'Connor I, O'Brien N. Modulation of UVA light-induced oxidative stress by β -carotene, lutein and astaxanthin in cultured fibroblasts. *J Dermatol Sci* 1998;16:226–30.
- [115] Fuchs J. Potentials and limitations of the natural antioxidants RRR- α -tocopherol, L-ascorbic acid and β -carotene in cutaneous photoprotection. *Free Radic Biol Med* 1998;25:848–73.
- [116] Anjani G, Ayustaningwarno F, Eviana R. Critical review on the immunomodulatory activities of carrot's β -carotene and other bioactive compounds. *J Funct Foods* 2022;99:105303.
- [117] Hughes DA. Effects of carotenoids on human immune function. *Proc Nutr Soc* 1999;58:713–8.
- [118] Hadad N, Levy R. The synergistic anti-inflammatory effects of lycopene, lutein, β -carotene, and carnosic acid combinations via redox-based inhibition of NF- κ B signalling. *Free Radic Biol Med* 2012;53:1381–91.
- [119] Wertz K, Hunziker PB, Seifert N, *et al.* β -carotene interferes with ultraviolet light A-induced gene expression by multiple pathways. *J Investigat Dermatol* 2005;124:428–34.
- [120] Furman-Toczek D, Zagórska-Dziok M, Dudra-Jastrzębska M, *et al.* A review of selected natural phytochemicals in preventing and treating malignant skin neoplasms. *J Pre Clin Clin Res* 2016;10:127–30.
- [121] Búfalo MC, Ferreira I, Costa G, *et al.* Propolis and its constituent caffeic acid suppress LPS-stimulated pro-inflammatory response by blocking NF- κ B and MAPK activation in macrophages. *J Ethnopharmacol* 2013;149:84–92.
- [122] Bharti AC, Aggarwal BB. Nuclear factor-kappa B and cancer: its role in prevention and therapy. *Biochem Pharmacol* 2002;64:883–8.
- [123] Dorai T, Aggarwal BB. Role of chemopreventive agents in cancer therapy. *Cancer Lett* 2004;215:129–40.
- [124] Desai SJ, Prickril B, Rasooly A. Mechanisms of phytonutrient modulation of cyclooxygenase-2 (COX-2) and inflammation related to cancer. *Nutr Cancer* 2018;70:350–75.
- [125] Guan C, Zhou X, Li H, *et al.* NF- κ B inhibitors gifted by nature: the anticancer promise of polyphenol compounds. *Biomed Pharmacother* 2022;156:113951.
- [126] Hashem S, Ali TA, Akhtar S, *et al.* Targeting cancer signalling pathways by natural products: exploring promising anti-cancer agents. *Biomed Pharmacother* 2022;150:113054.
- [127] Seeram NP. Berry fruits for cancer prevention: current status and future prospects. *J Agric Food Chem* 2008;56:630–5.
- [128] Scalbert A, Johnson IT, Saltmarsh M. Polyphenols: antioxidants and beyond. *Am J Clin Nutr* 2005;81:215S–7S.
- [129] de Kok TM, van Breda SG, Manson MM. Mechanisms of combined action of different chemopreventive dietary compounds: a review. *Eur J Nutr* 2008;47:51–9.
- [130] Sik B, Kapcsándi V, Székelyhidi R, *et al.* Recent advances in the analysis of rosmarinic acid from herbs in the *Lamiaceae* family. *Nat Prod Commun* 2019;14:1934578X19864216.
- [131] Nunes S, Madureira AR, Campos D, *et al.* Therapeutic and nutraceutical potential of rosmarinic acid—cytoprotective properties and pharmacokinetic profile. *Crit Rev Food Sci Nutr* 2017;57:1799–806.
- [132] Shekarchi M, Hajimehdipoor H, Saeidnia S, *et al.* Comparative study of rosmarinic acid content in some plants of *Labiatae* family. *Pharmacogn Mag* 2012;8:37–41.
- [133] Costa DC, Costa HS, Albuquerque TG, *et al.* Advances in phenolic compounds analysis of aromatic plants and their potential applications. *Trends Food Sci Technol* 2015;45:336–54.
- [134] Kumar P, Verma AK, Umaraw P, *et al.* Plant phenolics as natural preservatives in food system. *Plant Phenolics in Sustainable Agriculture*, Vol. 1. Springer; 2020. pp. 367–406.
- [135] Afaq F. Natural agents: cellular and molecular mechanisms of photoprotection. *Arch Biochem Biophys* 2011;508:144–51.
- [136] Pérez-Sánchez A, Barrajón-Catalán E, Herranz-López M, *et al.* Lemon balm extract (*Melissa officinalis*, L.) promotes melanogenesis and prevents UVB-induced oxidative stress and DNA damage in a skin cell model. *J Dermatol Sci* 2016;84:169–77.
- [137] Liu HM, Cheng MY, Xun MH, *et al.* Possible mechanisms of oxidative stress-induced skin cellular senescence, inflammation, and cancer and the therapeutic potential of plant polyphenols. *Int J Mol Sci* 2023;24:3755.
- [138] Rodríguez-Luna A, Ávila-Román J, Oliveira H, *et al.* Fucoxanthin and rosmarinic acid combination has anti-inflammatory effects through regulation of NLRP3 inflammasome in UVB-exposed HaCaT keratinocytes. *Mar Drugs* 2019;17:451.
- [139] Fernando PMDJ, Piao MJ, Kang KA, *et al.* Rosmarinic acid attenuates cell damage against UVB radiation-induced oxidative stress via enhancing antioxidant effects in human HaCaT cells. *Biomol Ther (Seoul)* 2016;24:75.
- [140] Sánchez-Campillo M, Gabaldon JA, Castillo J, *et al.* Rosmarinic acid, a photo-protective agent against UV and other ionizing radiations. *Food Chem Toxicol* 2009;47:386–92.
- [141] Pressi G, Bertaiola O, Guarnerio C, *et al.* In vitro cultured *Melissa officinalis* cells as effective ingredient to protect skin against oxidative stress, blue light, and infrared irradiations damages. *Cosmetics* 2021;8:23.

- [142] Han YH, Kee JY, Hong SH. Rosmarinic acid activates AMPK to inhibit metastasis of colorectal cancer. *Front Pharmacol* 2018;9:68.
- [143] Xu Y, Xu G, Liu L, *et al.* Anti-invasion effect of rosmarinic acid via the extracellular signal-regulated kinase and oxidation–reduction pathway in Ls174-T cells. *J Cell Biochem* 2010;111:370–9.
- [144] Lee C, Park GH, Ahn EM, *et al.* Protective effect of *Codium fragile* against UVB-induced pro-inflammatory and oxidative damages in HaCaT cells and BALB/c mice. *Fitoterapia* 2013;86:54–63.
- [145] Mengoni ES, Vichera G, Rigano LA, *et al.* Suppression of COX-2, IL-1 β and TNF- α expression and leukocyte infiltration in inflamed skin by bioactive compounds from *Rosmarinus officinalis* L. *Fitoterapia* 2011; 82:414–21.
- [146] da Silva GB, Manica D, da Silva AP, *et al.* Rosmarinic acid decreases viability, inhibits migration and modulates expression of apoptosis-related CASP8/CASP3/NLRP3 genes in human metastatic melanoma cells. *Chem Biol Interact* 2023;375:110427.
- [147] Zhao J, Xu L, Jin D, *et al.* Rosmarinic acid and related dietary supplements: potential applications in the prevention and treatment of cancer. *Biomolecules* 2022;12:1410.
- [148] Moore J, Yousef M, Tsiani E. Anticancer effects of rosemary (*Rosmarinus officinalis* L.) extract and rosemary extract polyphenols. *Nutrients* 2016;8:731.
- [149] Li H, Zhang Y, Chen HH, *et al.* Rosmarinic acid inhibits stem-like breast cancer through hedgehog and Bcl-2/Bax signalling pathways. *Pharmacogn Mag* 2019;15:600–6.
- [150] Jang YG, Hwang KA, Choi KC. Rosmarinic acid, a component of rosmarin tea, induced the cell cycle arrest and apoptosis through modulation of HDAC2 expression in prostate cancer cell lines. *Nutrients* 2018;10:1784.
- [151] da Silva GB, Yamauchi MA, Zanini D, *et al.* Novel possibility for cutaneous melanoma treatment by means of rosmarinic acid action on purinergic signalling. *Purinergic Signal* 2022;18:61–81.
- [152] Psotova J, Svobodova A, Kolarova H, *et al.* Photoprotective properties of *Prunella vulgaris* and rosmarinic acid on human keratinocytes. *J Photochem Photobiol B Biol* 2006;84:167–74.
- [153] Huang SS, Zheng RL. Rosmarinic acid inhibits angiogenesis and its mechanism of action in vitro. *Cancer Lett* 2006;239:271–80.
- [154] Gupta D, Sharma RR, Rashid H, *et al.* Rosmarinic acid alleviates ultraviolet mediated skin aging via attenuation of mitochondrial and ER stress responses. *Exp Dermatol* 2023;32:799–807.
- [155] Huang L, Chen J, Quan J, *et al.* Rosmarinic acid inhibits proliferation and migration, promotes apoptosis and enhances cisplatin sensitivity of melanoma cells through inhibiting ADAM17/EGFR/AKT/GSK3 β axis. *Bioengineered* 2021;12:3065–76.
- [156] Sharmila R, Ghosh K, Manoharan S. Rewiring of chemically modified MAPK pathway and its downstream targets by rosmarinic acid in mice dermal cancer. *J Inform Comput Sci* 2020;10:1047–64.
- [157] Lukmanul Hakkim F, Miura M, Matsuda N, *et al.* An in vitro evidence for caffeic acid, rosmarinic acid and trans cinnamic acid as a skin protectant against γ -radiation. *Int J Low Radiat* 2014;9:305–16.
- [158] Noor S, Mohammad T, Rub MA, *et al.* Biomedical features and therapeutic potential of rosmarinic acid. *Arch Pharm Res* 2022;45:205–28.
- [159] Esmeeta A, Adhikary S, Dharshnaa V, *et al.* Plant-derived bioactive compounds in colon cancer treatment: an updated review. *Biomed Pharmacother* 2022;153:113384.
- [160] Taheri JB, Azimi S, Rafeian N, *et al.* Herbs in dentistry. *Int Dent J* 2011; 61:287–96.
- [161] Veras KS, Fachel FNS, Teixeira HF, *et al.* Technological strategies applied for rosmarinic acid delivery through different routes – a review. *J Drug Deliv Sci Technol* 2022;68:103054.
- [162] Marchev AS, Vasileva LV, Amirova KM, *et al.* Rosmarinic acid – from bench to valuable applications in food industry. *Trends Food Sci Technol* 2021;117:182–93.
- [163] Swamy MK, Sinniah UR, Ghasemzadeh A. Anticancer potential of rosmarinic acid and its improved production through biotechnological interventions and functional genomics. *Appl Microbiol Biotechnol* 2018;102:7775–93.
- [164] Portz D, Koch E, Slusarenko AJ. Effects of garlic (*Allium sativum*) juice containing allicin on *Phytophthora infestans* and downy mildew of cucumber caused by *Pseudoperonospora cubensis*. *Eur J Plant Pathol* 2008;122:197–206.
- [165] Guíllamón E, Andreo-Martínez P, Mut-Salud N, *et al.* Beneficial effects of organosulfur compounds from *Allium cepa* on gut health: a systematic review. *Foods* 2021;10:1680.
- [166] Marchese A, Barbieri R, Sanches-Silva A, *et al.* Antifungal and antibacterial activities of allicin: a review. *Trends Food Sci Technol* 2016; 52:49–56.
- [167] Sbrana C, Avio L, Giovannetti M. Beneficial mycorrhizal symbionts affecting the production of health-promoting phytochemicals. *Electrophoresis* 2014;35:1535–46.
- [168] Touloupakis E, Ghanotakis DF. Nutraceutical use of garlic sulfur-containing compounds. *Adv Exp Med Biol* 2010;698:110–21.
- [169] Lawson LD, Hunsaker SM. Allicin bioavailability and bioequivalence from garlic supplements and garlic foods. *Nutrients* 2018;10:812.
- [170] Amagase H. Clarifying the real bioactive constituents of garlic. *J Nutr* 2006;136:716S–25S.
- [171] Wang HC, Pao J, Lin SY, *et al.* Molecular mechanisms of garlic-derived allyl sulfides in the inhibition of skin cancer progression. *Ann N Y Acad Sci* 2012;1271:44–52.
- [172] Jobani BM, Najafzadeh N, Mazani M, *et al.* Molecular mechanism and cytotoxicity of allicin and all-trans retinoic acid against CD44+ versus CD117+ melanoma cells. *Phytomedicine* 2018;48:161–9.
- [173] Omar SH, Al-Wabel NA. Organosulfur compounds and possible mechanism of garlic in cancer. *Saudi Pharm J* 2010;18:51–8.
- [174] Fatima N, Baqri SSR, Alsulimani A, *et al.* Phytochemicals from Indian ethnomedicines: promising prospects for the management of oxidative stress and cancer. *Antioxidants* 2021;10:1606.
- [175] Chahal DS, Sivamani RK, Rivkah Isseroff R, *et al.* Plant-based modulation of toll-like receptors: an emerging therapeutic model. *Phytother Res* 2013;27:1423–38.
- [176] Chu YL, Ho CT, Chung JG, *et al.* Allicin induces anti-human liver cancer cells through the p53 gene modulating apoptosis and autophagy. *J Agric Food Chem* 2013;61:9839–48.
- [177] Dubey SK, ed. *Natural Products and Nano-Formulations in Cancer Chemoprevention*. CRC Press; 2023.
- [178] Hashemy SI, Amiri H, Hosseini H, *et al.* PEGylated lecithin–chitosan–folic acid nanoparticles as nanocarriers of allicin for in vitro controlled release and anticancer effects. *Appl Biochem Biotechnol* 2023;195:4036–52.
- [179] Hasanpourghadi M, Yeng Looi C, Kumar Pandurangan A, *et al.* Phytometabolites targeting the Warburg effect in cancer cells: a mechanistic review. *Curr Drug Targets* 2017;18:1086–94.
- [180] Catanzaro E, Canistro D, Pellicioni V, *et al.* Anticancer potential of allicin: a review. *Pharmacol Res* 2022;177:106118.
- [181] Pandey MK, Gupta SC, Nabavizadeh A, *et al.* Regulation of cell signalling pathways by dietary agents for cancer prevention and treatment. *Semin Cancer Biol* 2017;46:158–81.
- [182] Hitl M, Kladar N, Gavarić N, *et al.* Garlic burn injuries – a systematic review of reported cases. *Am J Emerg Med* 2021;44:5–10.
- [183] Fariás-Campomanes AM, Horita CN, Pollonio MA, *et al.* Allicin-rich extract obtained from garlic by pressurized liquid extraction: quantitative determination of allicin in garlic samples. *Food Public Health* 2014;4:272–8.
- [184] Clarke JD, Dashwood RH, Ho E. Multi-targeted prevention of cancer by sulforaphane. *Cancer Lett* 2008;269:291–304.
- [185] Shapiro TA, Fahey JW, Wade KL, *et al.* Chemoprotective glucosinolates and isothiocyanates of broccoli sprouts: metabolism and excretion in humans. *Cancer Epidemiol Biomarkers Prevent* 2001;10:501–8.
- [186] Fahey JW, Zhang Y, Talalay P. Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc Natl Acad Sci U S A* 1997;94:10367–72.
- [187] Atwell LL, Hsu A, Wong CP, *et al.* Absorption and chemopreventive targets of sulforaphane in humans following consumption of broccoli sprouts or a myrosinase-treated broccoli sprout extract. *Mol Nutr Food Res* 2015;59:424–33.
- [188] Wang GC, Farnham M, Jeffery EH. Impact of thermal processing on sulforaphane yield from broccoli (*Brassica oleracea* L. ssp. italica). *J Agric Food Chem* 2012;60:6743–8.
- [189] Shokri S, Jegasothy H, Augustin MA, *et al.* Thermosonication for the production of sulforaphane rich broccoli ingredients. *Biomolecules* 2021;11:321.
- [190] Chinembiri TN, Du Plessis LH, Gerber M, *et al.* Review of natural compounds for potential skin cancer treatment. *Molecules* 2014;19: 11679–721.
- [191] Jiang X, Liu Y, Ma L, *et al.* Chemopreventive activity of sulforaphane. *Drug Des Devel Ther* 2018;12:2905–13.
- [192] Giudice A, Montella M. Activation of the Nrf2–ARE signalling pathway: a promising strategy in cancer prevention. *Bioessays* 2006;28: 169–81.

- [193] Reddy SA, Shelar SB, Dang T-M, *et al.* Sulforaphane and its methyl-carbonyl analogs inhibit the LPS-stimulated inflammatory response in human monocytes through modulating cytokine production, suppressing chemotactic migration and phagocytosis in a NF- κ B- and MAPK-dependent manner. *Int Immunopharmacol* 2015;24(no. 2):440–50.
- [194] Hamsa TP, Thejass P, Kuttan G. Induction of apoptosis by sulforaphane in highly metastatic B16F-10 melanoma cells. *Drug Chem Toxicol* 2011;34:332–40.
- [195] Bhat TA, Singh RP. Tumor angiogenesis – a potential target in cancer chemoprevention. *Food Chem Toxicol* 2008;46:1334–45.
- [196] Balasubramanian S, Chew YC, Eckert RL. Sulforaphane suppresses polycomb group protein level via a proteasome-dependent mechanism in skin cancer cells. *Mol Pharmacol* 2011;80:870–8.
- [197] Pop S, Enciu AM, Tarcomnicu I, *et al.* Phytochemicals in cancer prevention: modulating epigenetic alterations of DNA methylation. *Phytochem Rev* 2019;18:1005–24.
- [198] Park LK, Friso S, Choi SW. Nutritional influences on epigenetics and age-related disease. *Proc Nutr Soc* 2012;71:75–83.
- [199] Eom YS, Shah FH, Kim SJ. Sulforaphane induces cell differentiation, melanogenesis and also inhibit the proliferation of melanoma cells. *Eur J Pharmacol* 2022;921:174894.
- [200] Dickinson SE, Melton TF, Olson ER, *et al.* Inhibition of activator protein-1 by sulforaphane involves interaction with cysteine in the cFos DNA-binding domain: implications for chemoprevention of UVB-induced skin cancer. *Cancer Res* 2009;69:7103–10.
- [201] Houghton CA, Fassett RG, Coombes JS. Sulforaphane: translational research from laboratory bench to clinic. *Nutr Rev* 2013;71:709–26.
- [202] Cerdá B, Tomás-Barberán FA, Espín JC. Metabolism of antioxidant and chemopreventive ellagitannins from strawberries, raspberries, walnuts, and oak-aged wine in humans: identification of biomarkers and individual variability. *J Agric Food Chem* 2005;53:227–35.
- [203] Bakkalbaşı E, Menteş Ö, Artik N. Food ellagitannins – occurrence, effects of processing and storage. *Crit Rev Food Sci Nutr* 2008;49:283–98.
- [204] Hseu YC, Chou CW, Kumar KS, *et al.* Ellagic acid protects human keratinocyte (HaCaT) cells against UVA-induced oxidative stress and apoptosis through the upregulation of the HO-1 and Nrf-2 antioxidant genes. *Food Chem Toxicol* 2012;50:1245–55.
- [205] Devipriya N, Srinivasan M, Sudheer AR, *et al.* Effect of ellagic acid, a natural polyphenol, on alcohol-induced prooxidant and antioxidant imbalance: a drug dose dependent study. *Singapore Med J* 2007;48:311–8.
- [206] Umesalma S, Sudhandiran G. Differential inhibitory effects of the polyphenol ellagic acid on inflammatory mediators NF- κ B, iNOS, COX-2, TNF- α , and IL-6 in 1,2-dimethylhydrazine-induced rat colon carcinogenesis. *Basic Clin Pharmacol Toxicol* 2010;107:650–5.
- [207] Martin KR. Targeting apoptosis with dietary bioactive agents. *Exp Biol Med* 2006;231:117–29.
- [208] Malik A, Afaq S, Shahid M, *et al.* Influence of ellagic acid on prostate cancer cell proliferation: a caspase-dependent pathway. *Asian Pac J Trop Med* 2011;4:550–5.
- [209] D'Archivio M, Santangelo C, Scazzocchio B, *et al.* Modulatory effects of polyphenols on apoptosis induction: relevance for cancer prevention. *Int J Mol Sci* 2008;9:213–28.
- [210] Wang F, Chen J, Xiang D, *et al.* Ellagic acid inhibits cell proliferation, migration, and invasion in melanoma via EGFR pathway. *Am J Transl Res* 2020;12:2295.
- [211] Bai S, Yu Y, An L, *et al.* Ellagic acid increases stress resistance via insulin/IGF-1 signalling pathway in *Caenorhabditis elegans*. *Molecules* 2022;27:6168.
- [212] Zuccari G, Baldassari S, Ailuno G, *et al.* Formulation strategies to improve oral bioavailability of ellagic acid. *Appl Sci* 2020;10:3353.
- [213] Chatterjee S, Jain S, Jangid R, *et al.* Cytochrome P450 and P-gp mediated herb–drug interactions of some common Indian herbs. *Stud Nat Prod Chem* 2022;72:225–58.
- [214] Sharifi-Rad J, Quispe C, Castillo CMS, *et al.* Ellagic acid: a review on its natural sources, chemical stability, and therapeutic potential. *Oxid Med Cell Longev* 2022;2022:3848084.
- [215] González-Sarrías A, García-Villalba R, Núñez-Sánchez MÁ, *et al.* Identifying the limits for ellagic acid bioavailability: a crossover pharmacokinetic study in healthy volunteers after consumption of pomegranate extracts. *J Funct Foods* 2015;19:225–35.
- [216] Espín JC, García-Conesa MT, Tomás-Barberán FA. Nutraceuticals: facts and fiction. *Phytochemistry* 2007;68:2986–3008.
- [217] Hordyjewska A, Ostapiuk A, Horecka A, *et al.* Betulin and betulinic acid: triterpenoids derivatives with a powerful biological potential. *Phytochem Rev* 2019;18:929–51.
- [218] Mahajan RTCM, Chopda M. Phyto-pharmacology of *Ziziphus jujuba* Mill – a plant review. *Pharmacogn Rev* 2009;3:320.
- [219] Farneti B, Masuero D, Costa F, *et al.* Is there room for improving the nutraceutical composition of apple? *J Agric Food Chem* 2015;63:2750–9.
- [220] Ponou BK, Teponno RB, Taponjou AL, *et al.* Steroidal saponins from the aerial parts of *Cordylone fruticosa* L. var. strawberries. *Fitoterapia* 2019;134:454–8.
- [221] Demets OV, Takibayeva AT, Kassenov RZ, *et al.* Methods of betulin extraction from birch bark. *Molecules* 2022;27:3621.
- [222] Cichewicz RH, Kouzi SA. Chemistry, biological activity, and chemotherapeutic potential of betulinic acid for the prevention and treatment of cancer and HIV infection. *Med Res Rev* 2004;24:90–114.
- [223] Ríos JL, Máñez S. New pharmacological opportunities for betulinic acid. *Planta Med* 2018;84:8–19.
- [224] Tewari D, Patni P, Bishayee A, *et al.* Natural products targeting the PI3K-Akt-mTOR signalling pathway in cancer: a novel therapeutic strategy. *Semin Cancer Biol* 2022;30:1–17.
- [225] Gangwar V, Garg A, Lomore K, *et al.* Immunomodulatory effects of a concoction of natural bioactive compounds—mechanistic insights. *Biomedicines* 2021;9:1522.
- [226] Sugiura R, Satoh R, Takasaki T. ERK: a double-edged sword in cancer. ERK-dependent apoptosis as a potential therapeutic strategy for cancer. *Cells* 2021;10:2509.
- [227] Sardar A, Gautam S, Sinha S, *et al.* Nanoparticles of naturally occurring PPAR- γ inhibitor betulinic acid ameliorates bone marrow adiposity and pathological bone loss in ovariectomized rats via Wnt/ β -catenin pathway. *Life Sci* 2022;309:121020.
- [228] Wróblewska-Luczka P, Cabaj J, Bąk W, *et al.* Additive interactions between betulinic acid and two taxanes in in vitro tests against four human malignant melanoma cell lines. *Int J Mol Sci* 2022;23:9641.
- [229] Ghiulai R, Mioc A, Racoviceanu R, *et al.* The anti-melanoma effect of betulinic acid functionalized gold nanoparticles: a mechanistic in vitro approach. *Pharmaceutics* 2022;15:1362.
- [230] Kallimanis P, Chinou I, Panagiotopoulou A, *et al.* *Rosmarinus officinalis* L. leaf extracts and their metabolites inhibit the aryl hydrocarbon receptor (AhR) activation in vitro and in human keratinocytes: potential impact on inflammatory skin diseases and skin cancer. *Molecules* 2022;27:2499.
- [231] Bravo-Alfaro DA, Ochoa-Rodríguez LR, Villaseñor-Ortega F, *et al.* Self-nanoemulsifying drug delivery system (SNEDDS) improves the oral bioavailability of betulinic acid. *J Mol Liquids* 2022;364:119946.
- [232] Ameta RK, Soni K, Bhattarai A. Recent advances in improving the bioavailability of hydrophobic/lipophilic drugs and their delivery via self-emulsifying formulations. *Colloids Interfaces* 2023;7:16.
- [233] Wu X, Wei Z, Feng H, *et al.* Targeting effect of betulinic acid liposome modified by hyaluronic acid on hepatoma cells in vitro. *J Pharm Sci* 2022;111:3047–53.
- [234] Peng F, Jin Y, Wang K, *et al.* Glycosylated zein composite nanoparticles for efficient delivery of betulinic acid: fabrication, characterization, and in vitro release properties. *Foods* 2022;11:2589.
- [235] Kumar V, Garg V, Dureja H. Nanoemulsion for delivery of anticancer drugs. *Cancer Adv* 2022;5:e22016.
- [236] Yogeeswari P, Sriram D. Betulinic acid and its derivatives: a review on their biological properties. *Curr Med Chem* 2005;12:657–66.
- [237] Shankar E, Goel A, Gupta K, *et al.* Plant flavone apigenin: an emerging anticancer agent. *Curr Pharmacol Rep* 2017;3:423–46.
- [238] Wang H, Yang L, Zu Y, *et al.* Microwave-assisted simultaneous extraction of luteolin and apigenin from tree peony pod and evaluation of its antioxidant activity. *ScientificWorldJournal* 2014;2014:506971.
- [239] Michalaki A, Karantonis HC, Kritikou AS, *et al.* Ultrasound-assisted extraction of specific phenolic compounds from petroselinum crispum leaves using response surface methodology and HPLC-PDA and Q-TOF-MS/MS identification. *Appl Sci* 2023;13:798.
- [240] Saotome Y, Imai M. Supercritical carbon dioxide extraction of apigenin from parsley leaves pre-treated to maximize yield. *Food Sci Technol Res* 2018;24:63–73.
- [241] Wang R, Ni N, Zhang QX, *et al.* Adsorption and separation of flavonoid aglycones and flavonol aglycones by using Zr (IV) immobilized collagen fiber adsorbent as column packing material. *Sep Purif Technol* 2022;289:120684.

- [242] Parihar A, Eubank TD, Doseff AI. Monocytes and macrophages regulate immunity through dynamic networks of survival and cell death. *J Innate Immun* 2010;2:204–15.
- [243] Aggarwal BB, Shishodia S. Molecular targets of dietary agents for prevention and therapy of cancer. *Biochem Pharmacol* 2006;71:1397–421.
- [244] Ozbey U, Attar R, Romero MA, et al. Apigenin as an effective anticancer natural product: spotlight on TRAIL, WNT/ β -catenin, JAK-STAT pathways, and microRNAs. *J Cell Biochem* 2019;120:1060–7.
- [245] L Suraweera T, Rupasinghe HV, Dellaire G, et al. Regulation of Nrf2/ARE pathway by dietary flavonoids: a friend or foe for cancer management? *Antioxidants* 2020;9:973.
- [246] Montes de Oca MK, Pearlman RL, McClees SF, et al. Phytochemicals for the prevention of photocarcinogenesis. *Photochem Photobiol* 2017;93:956–74.
- [247] Bridgeman BB, Wang P, Ye B, et al. Inhibition of mTOR by apigenin in UVB-irradiated keratinocytes: a new implication of skin cancer prevention. *Cell Signal* 2016;28:460–8.
- [248] Das S, Das J, Samadder A, et al. Strategic formulation of apigenin-loaded PLGA nanoparticles for intracellular trafficking, DNA targeting and improved therapeutic effects in skin melanoma in vitro. *Toxicol Lett* 2013;223:124–38.
- [249] Jangdey MS, Gupta A, Saraf S, et al. Development and optimization of apigenin-loaded transdermal system for skin cancer delivery: in vitro evaluation. *Artif Cells Nanomed Biotechnol* 2017;45:1452–62.
- [250] Waheed A, Zameer S, Sultana N, et al. Engineering of QbD driven and ultrasonically shaped lyotropic liquid crystalline nanoparticles for Apigenin in the management of skin cancer. *Eur J Pharm Biopharm* 2022;180:269–80.
- [251] Fernández J, Silván B, Entrialgo-Cadierno R, et al. Antiproliferative and palliative activity of flavonoids in colorectal cancer. *Biomed Pharmacother* 2021;143:112241.
- [252] Ali F, Rahul, Naz F, et al. Health functionality of apigenin: a review. *Int J Food Prop* 2017;20:1197–238.
- [253] Zhou X, Fu L, Wang P, et al. Drug-herb interactions between *Scutellaria baicalensis* and pharmaceutical drugs: insights from experimental studies, mechanistic actions to clinical applications. *Biomed Pharmacother* 2021;138:111445.
- [254] Tang D, Chen K, Huang L, et al. Pharmacokinetic properties and drug interactions of apigenin, a natural flavone. *Exp Opin Drug Metab Toxicol* 2017;13:323–30.
- [255] Shahrajabian MH, Sun W, Cheng Q. Clinical aspects and health benefits of ginger (*Zingiber officinale*) in both traditional Chinese medicine and modern industry. *Acta Agric Scand B Soil Plant Sci* 2019;69:546–56.
- [256] Cosme P, Rodríguez AB, Espino J, et al. Plant phenolics: bioavailability as a key determinant of their potential health-promoting applications. *Antioxidants* 2020;9:1263.
- [257] Semwal RB, Semwal DK, Combrinck S, et al. Gingerols and shogaols: important nutraceutical principles from ginger. *Phytochemistry* 2015;117:554–68.
- [258] Ali BH, Blunden G, Tanira MO, et al. Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food Chem Toxicol* 2008;46:409–20.
- [259] Nafees S, Zafaryab M, Mehdi SH, et al. Anti-cancer effect of gingerol in cancer prevention and treatment. *Anticancer Agents Med Chem* 2021;21:428–32.
- [260] Anbar EA, El-Naggar SA, Salama AF. The rationale of gingerol as a main phenolic compound of ginger in improving hepatocellular carcinoma induced by dimethylaminoazobenzene: scientific implication. *Egypt Acad J Biol Sci D Histol Histochem* 2022;14:25–30.
- [261] White B. Ginger: an overview. *Am Fam Physician* 2007;75:1689–91.
- [262] Sharifi-Rad M, Varoni EM, Salehi B, et al. Plants of the genus *Zingiber* as a source of bioactive phytochemicals: from tradition to pharmacy. *Molecules* 2017;22:2145.
- [263] Surh YJ. Cancer chemoprevention with dietary phytochemicals. *Nat Rev Cancer* 2003;3:768–80.
- [264] Park KK, Chun KS, Lee JM, et al. Inhibitory effects of [6]-gingerol, a major pungent principle of ginger, on phorbol ester-induced inflammation, epidermal ornithine decarboxylase activity and skin tumor promotion in ICR mice. *Cancer Lett* 1998;129:139–44.
- [265] Nigam N, George J, Srivastava S, et al. Induction of apoptosis by [6]-gingerol associated with the modulation of p53 and involvement of mitochondrial signalling pathway in B[a]P-induced mouse skin tumorigenesis. *Cancer Chemother Pharmacol* 2010;65:687–96.
- [266] Adikesavan M, Athiraja P, Divakar MBB. Investigation on the anticancer activity of [6]-gingerol of *Zingiber officinale* and its structural analogs against skin cancer. *Curr Comput Aided Drug Des* 2024;20:367–73.
- [267] Abebe W. Review of herbal medications with the potential to cause bleeding: dental implications, and risk prediction and prevention avenues. *EPMA J* 2019;10:51–64.
- [268] Alipour A, Baradaran Rahimi V, Askari VR. Promising influences of gingerols against metabolic syndrome: a mechanistic review. *Biofactors* 2022;48:993–1004.
- [269] Tiran D. Ginger to reduce nausea and vomiting during pregnancy: evidence of effectiveness is not the same as proof of safety. *Complement Ther Clin Pract* 2012;18:22–5.
- [270] Iqbal MK, Chaudhuri A, Iqbal A, et al. Targeted delivery of natural bioactives and lipid-nanocargos against signalling pathways involved in skin cancer. *Curr Med Chem* 2021;28:8003–35.
- [271] Lakhanpal P, Rai DK. Quercetin: a versatile flavonoid. *Internet J Med Update* 2007;2:22–37.
- [272] Slimestad R, Fossen T, Vågen IM. Onions: a source of unique dietary flavonoids. *J Agric Food Chem* 2007;55:10067–80.
- [273] Boyer J, Liu RH. Apple phytochemicals and their health benefits. *Nutr J* 2004;3:5.
- [274] Seeram NP. Berry fruits: compositional elements, biochemical activities, and the impact of their intake on human health, performance, and disease. *J Agric Food Chem* 2008;56:627–9.
- [275] Franke AA, Custer LJ, Arakaki C, et al. Vitamin C and flavonoid levels of fruits and vegetables consumed in Hawaii. *J Food Compos Anal* 2004;17:1–35.
- [276] Jaganath IB, Crozier A. Dietary flavonoids and phenolic compounds. *Plant Phenolics and Human Health: Biochemistry, Nutrition, and Pharmacology*, 2010. pp. 1–49.
- [277] Ivana B, Marija L. Role of phenols in energy and functional beverages. *Sports and Energy Drinks*, 2019. pp. 229–268.
- [278] Lesjak M, Beara I, Simin N, et al. Antioxidant and anti-inflammatory activities of quercetin and its derivatives. *J Funct Foods* 2018;40:68–75.
- [279] Bądziul D, Jakubowicz-Gil J, Langner E, et al. The effect of quercetin and imperatorin on programmed cell death induction in T98G cells in vitro. *Pharmacol Rep* 2014;66:292–300.
- [280] Dhillon AS, Hagan S, Rath O, et al. MAP kinase signalling pathways in cancer. *Oncogene* 2007;26:3279–90.
- [281] Anastas JN, Moon RT. WNT signalling pathways as therapeutic targets in cancer. *Nat Rev Cancer* 2013;13:11–26.
- [282] Li H, Chen C. Quercetin has antimetastatic effects on gastric cancer cells via the interruption of uPA/uPAR function by modulating NF- κ B, PKC- δ , ERK1/2, and AMPK α . *Integr Cancer Ther* 2018;17:511–23.
- [283] Imran M, Iqbal MK, Imtiyaz K, et al. Topical nanostructured lipid carrier gel of quercetin and resveratrol: formulation, optimization, in vitro and ex vivo study for the treatment of skin cancer. *Int J Pharm* 2020;587:119705.
- [284] Caddeo C, Nacher A, Vassallo A, et al. Effect of quercetin and resveratrol co-incorporated in liposomes against inflammatory/oxidative response associated with skin cancer. *Int J Pharm* 2016;513:153–63.
- [285] Paliwal S, Sundaram J, Mitragotri S. Induction of cancer-specific cytotoxicity towards human prostate and skin cells using quercetin and ultrasound. *Br J Cancer* 2005;92:499–502.
- [286] Jung M, Bu SY, Tak KH, et al. Anticarcinogenic effect of quercetin by inhibition of insulin-like growth factor (IGF)-1 signalling in mouse skin cancer. *Nutr Res Pract* 2013;7:439–45.
- [287] D'Andrea G. Quercetin: a flavonol with multifaceted therapeutic applications? *Fitoterapia* 2015;106:256–71.
- [288] Di Bari L, Ripoli S, Pradhan S, et al. Interactions between quercetin and warfarin for albumin binding: a new eye on food/drug interference. *Chirality* 2010;22:593–6.
- [289] Samuel T, Fadlalla K, Mosley L, et al. Dual-mode interaction between quercetin and DNA-damaging drugs in cancer cells. *Anticancer Res* 2012;32:61–71.
- [290] Gupta A, Birhman K, Raheja I, et al. Quercetin: a wonder bioflavonoid with therapeutic potential in disease management. *Asian Pac J Trop Dis* 2016;6:248–52.
- [291] Simoes MF, Sousa JS, Pais AC. Skin cancer and new treatment perspectives: a review. *Cancer Lett* 2015;357:8–42.
- [292] Montané X, Kowalczyk O, Reig-Vano B, et al. Current perspectives of the applications of polyphenols and flavonoids in cancer therapy. *Molecules* 2020;25:3342.

- [293] Liu Y, Cai X, Zhang Y, *et al.* Kaempferol alleviates renal fibrosis via NF- κ B-mediated inhibition of epithelial-to-mesenchymal transition in vitro and in vivo. *Food Fun* 2020;11:2972–85.
- [294] Xu W, Liu J, Li C, *et al.* Kaempferol-7-O- β -D-glucoside (KG) isolated from *Smilax china* L. rhizome induces G2/M phase arrest and apoptosis on HeLa cells in a p53-independent manner. *Cancer Lett* 2008;264:229–40.
- [295] Lee HS, Cho HJ, Yu R, *et al.* Mechanisms underlying apoptosis-inducing effects of Kaempferol in HT-29 human colon cancer cells. *Int J Mol Sci* 2014;15:2722–37.
- [296] Nam SY, Jeong HJ, Kim HM. Kaempferol impedes IL-32-induced monocyte-macrophage differentiation. *Chem Biol Interact* 2017;274:107–15.
- [297] Liao W, Chen L, Ma X, *et al.* Protective effects of kaempferol against reactive oxygen species-induced hemolysis and its antiproliferative activity on human cancer cells. *Eur J Med Chem* 2016;114:24–32.
- [298] Farasat A, Ghorbani M, Gheibi N, *et al.* In silico assessment of the inhibitory effect of four flavonoids (Chrysin, Naringin, Quercetin, Kaempferol) on tyrosinase activity using the MD simulation approach. *BioTechnologia J Biotechnol Comput Biol Bionanotechnol* 2020;101:193–204.
- [299] Li C, Zhao Y, Yang D, *et al.* Inhibitory effects of kaempferol on the invasion of human breast carcinoma cells by downregulating the expression and activity of matrix metalloproteinase-9. *Biochem Cell Biol* 2015;93:16–27.
- [300] Rajendran P, Rengarajan T, Nandakumar N, *et al.* Kaempferol, a potential cytostatic and cure for inflammatory disorders. *Eur J Med Chem* 2014;86:103–12.
- [301] Yang J, Xiao P, Sun J, *et al.* Anticancer effects of kaempferol in A375 human malignant melanoma cells are mediated via induction of apoptosis, cell cycle arrest, inhibition of cell migration and downregulation of m-TOR/PI3K/AKT pathway. *J BUON* 2018;23:218–23.
- [302] Yao K, Chen H, Liu K, *et al.* Kaempferol targets RSK2 and MSK1 to suppress UV radiation-induced skin cancer. *Cancer Prev Res (Phila)* 2014;7:958–67.
- [303] Lee KM, Lee KW, Jung SK, *et al.* Kaempferol inhibits UVB-induced COX-2 expression by suppressing Src kinase activity. *Biochem Pharmacol* 2010;80:2042–9.
- [304] Kim SH, Choi KC. Anti-cancer effect and underlying mechanism(s) of kaempferol, a phytoestrogen, on the regulation of apoptosis in diverse cancer cell models. *Toxicol Res* 2013;29:229–34.
- [305] Dabeek WM, Marra MV. Dietary quercetin and kaempferol: bioavailability and potential cardiovascular-related bioactivity in humans. *Nutrients* 2019;11:2288.
- [306] Lambert JD, Sang S, Yang CS. Possible controversy over dietary polyphenols: benefits vs risks. *Chem Res Toxicol* 2007;20:583–5.
- [307] Hodek P, Trefil P, Stiborová M. Flavonoids-potent and versatile biologically active compounds interacting with cytochromes P450. *Chem Biol Interact* 2002;139:1–21.
- [308] Igura K, Ohta T, Kuroda Y, *et al.* Resveratrol and quercetin inhibit angiogenesis in vitro. *Cancer Lett* 2001;171:11–6.
- [309] Bouby L, Figueiral I, Bouchette A, *et al.* Bioarchaeological insights into the process of domestication of grapevine (*Vitis vinifera* L.) during Roman times in Southern France. *PLoS One* 2013;8:e63195.
- [310] Moreno A, Castro M, Falqué E. Evolution of trans-and cis-resveratrol content in red grapes (*Vitis vinifera* L. cv Mencía, Albarello and Merenzao) during ripening. *Eur Food Res Technol* 2008;227:667–74.
- [311] Sales JM, Resurreccion AV. Resveratrol in peanuts. *Crit Rev Food Sci Nutr* 2014;54:734–70.
- [312] Ector BJ, Magee JB, Hegwood CP, *et al.* Resveratrol concentration in muscadine berries, juice, pomace, purees, seeds, and wines. *Am J Enol Vitic* 1996;47:57–62.
- [313] Chen H, Tuck T, Ji X, *et al.* Quality assessment of Japanese knotweed (*Fallopia japonica*) grown on Prince Edward Island as a source of resveratrol. *J Agric Food Chem* 2013;61:6383–92.
- [314] Charles DJ. Sources of natural antioxidants and their activities. *Antioxidant Properties of Spices, Herbs and Other Sources*, 2013. pp. 65–138.
- [315] Jeandet P, Bessis R, Sbaghi M, *et al.* Production of the phytoalexin resveratrol by grapes as a response to Botrytis attack under natural conditions. *J Phytopathol* 1995;143:135–9.
- [316] Kuršvietienė L, Stanevičienė I, Mongirdienė A, *et al.* Multiplicity of effects and health benefits of resveratrol. *Medicina* 2016;52:148–55.
- [317] Švajger U, Jeras M. Anti-inflammatory effects of resveratrol and its potential use in therapy of immune-mediated diseases. *Int Rev Immunol* 2012;31:202–22.
- [318] Dybkowska E, Sadowska A, Swiderski F, *et al.* The occurrence of resveratrol in foodstuffs and its potential for supporting cancer prevention and treatment. A review. *Rocz Państw Zakł Hig* 2018;69:5–14.
- [319] Lee IT, Yang CM. Role of NADPH oxidase/ROS in pro-inflammatory mediators-induced airway and pulmonary diseases. *Biochem Pharmacol* 2012;84:581–90.
- [320] Kundu JK, Surh YJ. Inflammation: gearing the journey to cancer. *Mutat Res* 2008;659:15–30.
- [321] Zhao H, Han L, Jian Y, *et al.* Resveratrol induces apoptosis in human melanoma cell through negatively regulating Erk/PKM2/Bcl-2 axis. *Oncotargets Ther* 2018;11:8995–9006.
- [322] Gali-Muhtasib H, Hmadi R, Kareh M, *et al.* Cell death mechanisms of plant-derived anticancer drugs: beyond apoptosis. *Apoptosis* 2015;20:1531–62.
- [323] Yan Y, Yang JY, Mou YH, *et al.* Differences in the activities of resveratrol and ascorbic acid in protection of ethanol-induced oxidative DNA damage in human peripheral lymphocytes. *Food Chem Toxicol* 2012;50:168–74.
- [324] Pratheeshkumar P, Sreekala C, Zhang Z, *et al.* Cancer prevention with promising natural products: mechanisms of action and molecular targets. *Anticancer Agents Med Chem* 2012;12:1159–84.
- [325] Osmond GW, Augustine CK, Zipfel PA, *et al.* Enhancing melanoma treatment with resveratrol. *J Surg Res* 2012;172:109–15.
- [326] Aziz SW, Aziz MH. Protective molecular mechanisms of resveratrol in UVR-induced skin carcinogenesis. *Photodermatol Photoimmunol Photomed* 2018;34:35–41.
- [327] Berman AY, Motechin RA, Wiesenfeld MY, *et al.* The therapeutic potential of resveratrol: a review of clinical trials. *NPJ Precis Oncol* 2017;1:35.
- [328] Kundu JK, Shin YK, Kim SH, *et al.* Resveratrol inhibits phorbol ester-induced expression of COX-2 and activation of NF- κ B in mouse skin by blocking I κ B kinase activity. *Carcinogenesis* 2006;27:1465–74.
- [329] Reagan-Shaw S, Afaq F, Aziz MH, *et al.* Modulations of critical cell cycle regulatory events during chemoprevention of ultraviolet B-mediated responses by resveratrol in SKH-1 hairless mouse skin. *Oncogene* 2004;23:5151–60.
- [330] Wu Z, Uchi H, Morino-Koga S, *et al.* Resveratrol inhibition of human keratinocyte proliferation via SIRT1/ARNT/ERK dependent down-regulation of aquaporin 3. *J Dermatolog Sci* 2014;75:16–23.
- [331] Chiba T, Kimura Y, Suzuki S, *et al.* Trans-resveratrol enhances the anticoagulant activity of warfarin in a mouse model. *J Atheroscler Thromb* 2016;23:1099–110.
- [332] Gescher A. Polyphenolic phytochemicals versus non-steroidal anti-inflammatory drugs: which are better cancer chemopreventive agents? *J Chemother* 2004;16:3–6.
- [333] Lu C, Lin HZ, Li YY, *et al.* Making room for specific molecules in the treatment of depression. *Sci J Depress Anxiety* 2017;1:20–2.
- [334] Jiang Z, Chen K, Cheng L, *et al.* Resveratrol and cancer treatment: updates. *Ann N Y Acad Sci* 2017;1403:59–69.
- [335] Ren B, Kwah MX-Y, Liu C, *et al.* Resveratrol for cancer therapy: challenges and future perspectives. *Cancer Lett* 2021;515:63–72.
- [336] Chedea VS, Vicaș SI, Sticozzi C, *et al.* Resveratrol: from diet to topical usage. *Food Funct* 2017;8:3879–92.
- [337] Fibigr J, Šatinský D, Solich P. Current trends in the analysis and quality control of food supplements based on plant extracts. *Anal Chim Acta* 2018;1036:1–15.
- [338] Akram M, Shahab-Uddin AA, Usmanghani K, *et al.* Curcuma longa and curcumin: a review article. *Rom J Biol Plant Biol* 2010;55:65–70.
- [339] Debjit Bhowmik C, Kumar KS, Chandira M, *et al.* Turmeric: a herbal and traditional medicine. *Arch Appl Sci Res* 2009;1:86–108.
- [340] Abd El-Hack ME, El-Saadony MT, Swelum AA, *et al.* Curcumin, the active substance of turmeric: its effects on health and ways to improve its bioavailability. *J Sci Food Agric* 2021;101:5747–62.
- [341] Ak T, Gülçin I. Antioxidant and radical scavenging properties of curcumin. *Chem Biol Interact* 2008;174:27–37.
- [342] Menon VP and Sudheer AR. Antioxidant and anti-inflammatory properties of curcumin. In: Aggarwal BB, Surh Y-J, Shishodia S, eds. *The Molecular Targets and Therapeutic Uses of Curcumin in Health and Disease*. Springer; 2007. pp. 105–25.

- [343] Perrone D, Ardito F, Giannatempo G, *et al.* Biological and therapeutic activities, and anticancer properties of curcumin. *Exp Ther Med* 2015; 10:1615–23.
- [344] Jobin C, Bradham CA, Russo MP, *et al.* Curcumin blocks cytokine-mediated NF- κ B activation and proinflammatory gene expression by inhibiting inhibitory factor I- κ B kinase activity. *J Immunol* 1999;163:3474–83.
- [345] Reuter S, Eifes S, Dicato M, *et al.* Modulation of anti-apoptotic and survival pathways by curcumin as a strategy to induce apoptosis in cancer cells. *Biochem Pharmacol* 2008;76:1340–51.
- [346] Zhao G, Han X, Zheng S, *et al.* Curcumin induces autophagy, inhibits proliferation and invasion by downregulating AKT/mTOR signalling pathway in human melanoma cells. *Oncol Rep* 2016;35:1065–74.
- [347] Wu J, Lu WY, Cui LL. Inhibitory effect of curcumin on invasion of skin squamous cell carcinoma A431 cells. *Asian Pac J Cancer Prev* 2015;16: 2813–8.
- [348] Kunnumakkara AB, Diagaradjane P, Anand P, *et al.* Curcumin sensitizes human colorectal cancer xenografts in nude mice to gamma-radiation by targeting nuclear factor-kappaB-regulated gene products. *Clin Cancer Res* 2008;14:2128–36.
- [349] Rein MJ, Renouf M, Cruz-Hernandez C, *et al.* Bioavailability of bioactive food compounds: a challenging journey to bioefficacy. *Br J Clin Pharmacol* 2013;75:588–602.
- [350] Wang Z, Sun W, Huang CK, *et al.* Inhibitory effects of curcumin on activity of cytochrome P450 2C9 enzyme in human and 2C11 in rat liver microsomes. *Drug Dev Ind Pharm* 2015;41:613–6.
- [351] Hu C, Kitts DD. Evaluation of antioxidant activity of epigallocatechin gallate in biphasic model systems in vitro. *Mol Cell Biochem* 2001;218: 147–55.
- [352] Zhong Y, Chiou YS, Pan MH, *et al.* Anti-inflammatory activity of lipophilic epigallocatechin gallate (EGCG) derivatives in LPS-stimulated murine macrophages. *Food Chem* 2012;134:742–8.
- [353] Namita P, Mukesh R, Vijay KJ. *Camellia sinensis* (green tea): a review. *Glob J Pharmacol* 2012;6:52–9.
- [354] Łuczaj W, Skrzydlewska E. Antioxidative properties of black tea. *Prev Med* 2005;40:910–8.
- [355] Pastoriza S, Mesías M, Cabrera C, *et al.* Healthy properties of green and white teas: an update. *Food Funct* 2017;8:2650–62.
- [356] Lee RJ, Lee VS, Tzen JT, *et al.* Study of the release of gallic acid from (–)-epigallocatechin gallate in old oolong tea by mass spectrometry. *Rapid Commun Mass Spectrom* 2010;24:851–8.
- [357] Ferrari E, Bettuzzi S, Naponelli V. The potential of epigallocatechin gallate (EGCG) in targeting autophagy for cancer treatment: a narrative review. *Int J Mol Sci* 2022;23:6075.
- [358] Eng QY, Thanikachalam PV, Ramamurthy S. Molecular understanding of epigallocatechin gallate (EGCG) in cardiovascular and metabolic diseases. *J Ethnopharmacol* 2018;210:296–310.
- [359] Wightman EL, Haskell CF, Forster JS, *et al.* Epigallocatechin gallate, cerebral blood flow parameters, cognitive performance and mood in healthy humans: a double-blind, placebo-controlled, crossover investigation. *Hum Psychopharmacol* 2012;27:177–86.
- [360] Oz HS. Chronic inflammatory diseases and green tea polyphenols. *Nutrients* 2017;9:561.
- [361] H Farzaei M, Rahimi R, Abdollahi M. The role of dietary polyphenols in the management of inflammatory bowel disease. *Curr Pharm Biotechnol* 2015;16:196–210.
- [362] Liu Q, Qian Y, Chen F, *et al.* EGCG attenuates pro-inflammatory cytokines and chemokines production in LPS-stimulated L02 hepatocyte. *Acta Biochim Biophys Sin (Shanghai)* 2014;46:31–9.
- [363] Gan R-Y, Li H-B, Sui Z-Q, *et al.* Absorption, metabolism, anti-cancer effect and molecular targets of epigallocatechin gallate (EGCG): an updated review. *Crit Rev Food Sci Nutr* 2018;58:924–41.
- [364] El-Kayal M, Nasr M, Elkheshen S, *et al.* Colloidal (–)-epigallocatechin-3-gallate vesicular systems for prevention and treatment of skin cancer: a comprehensive experimental study with preclinical investigation. *Eur J Pharm Sci* 2019;137:104972.
- [365] Zhang J, Lei Z, Huang Z, *et al.* Epigallocatechin-3-gallate (EGCG) suppresses melanoma cell growth and metastasis by targeting TRAF6 activity. *Oncotarget* 2016;7:79557.
- [366] Ellis LZ, Liu W, Luo Y, *et al.* Green tea polyphenol epigallocatechin-3-gallate suppresses melanoma growth by inhibiting inflammasome and IL-1 β secretion. *Biochem Biophys Res Commun* 2011;414:551–6.
- [367] Lambert JD, Kennett MJ, Sang S, *et al.* Hepatotoxicity of high oral dose (–)-epigallocatechin-3-gallate in mice. *Food Chem Toxicol* 2010;48: 409–16.
- [368] Yang CS, Wang X, Lu G, *et al.* Cancer prevention by tea: animal studies, molecular mechanisms and human relevance. *Nat Rev Cancer* 2009;9: 429–39.
- [369] Andreu Fernandez V, Almeida Toledano L, Pizarro Lozano N, *et al.* Bioavailability of epigallocatechin gallate administered with different nutritional strategies in healthy volunteers. *Antioxidants* 2020;9:440.
- [370] Harati K, Behr B, Wallner C, *et al.* Anti-proliferative activity of epigallocatechin-3-gallate and silibinin on soft tissue sarcoma cells. *Mol Med Rep* 2017;15:103–10.
- [371] Zhang Y, Owusu L, Duan W, *et al.* Anti-metastatic and differential effects on protein expression of epigallocatechin-3-gallate in HCCLM6 hepatocellular carcinoma cells. *Int J Mol Med* 2013;32:959–64.
- [372] Guzman I, Hamby S, Romero J, *et al.* Variability of carotenoid biosynthesis in orange colored *Capsicum* spp. *Plant Sci* 2010;179:49–59.
- [373] Olson JA. Provitamin A function of carotenoids: the conversion of β -carotene into vitamin A. *J Nutr* 1989;119:105–8.
- [374] Johnson EJ. The role of carotenoids in human health. *Nutr Clin Care* 2002;5:56–65.
- [375] Gul K, Tak A, Singh AK, *et al.* Chemistry, encapsulation, and health benefits of β -carotene - a review. *Cogent Food Agric* 2015;1:1018696.
- [376] Wood SM, Beckham C, Yosioka A, *et al.* β -Carotene and selenium supplementation enhances immune response in aged humans. *Integr Med* 2000;2:85–92.
- [377] Pritwani R, Mathur P. β -carotene content of some commonly consumed vegetables and fruits available in Delhi, India. *J Nutr Food Sci* 2017;7: 1–7.
- [378] Devasagayam TPA, Tilak JC, Boloor KK, *et al.* Free radicals and antioxidants in human health: current status and future prospects. *J Assoc Physicians India* 2004;52:794–804.
- [379] Godic A, Poljšak B, Adamic M, *et al.* The role of antioxidants in skin cancer prevention and treatment. *Oxid Med Cell Longev* 2014;2014: 860479.
- [380] Greenberg ER, Baron JA, Stukel TA, *et al.* A clinical trial of beta carotene to prevent basal-cell and squamous-cell cancers of the skin. *N Engl J Med* 1990;323:789–95.
- [381] Kune GA, Bannerman S, Field B, *et al.* Diet, alcohol, smoking, serum β -carotene, and vitamin A in male nonmelanocytic skin cancer patients and controls. *Nutr Cancer* 1992;18:237–44.
- [382] Black HS, Boehm F, Edge R, *et al.* The benefits and risks of certain dietary carotenoids that exhibit both anti- and pro-oxidative mechanisms - a comprehensive review. *Antioxidants (Basel)* 2020;9:264.
- [383] Stahl W, Sies H. β -Carotene and other carotenoids in protection from sunlight. *Am J Clin Nutr* 2012;96:1179S–84S.
- [384] Seifried HE, McDonald SS, Anderson DE, *et al.* The antioxidant condurum in cancer. *Cancer Res* 2003;63:429S–8.
- [385] Tsai J, Chien AL. Photoprotection for skin of color. *Am J Clin Dermatol* 2022;23:195–205.
- [386] Alam MA, Subhan N, Hossain H, *et al.* Hydroxycinnamic acid derivatives: a potential class of natural compounds for the management of lipid metabolism and obesity. *Nutr Metab* 2016;13:1–13.
- [387] Crozier A, Yokota T, Jaganath IB, *et al.* Secondary metabolites in fruits, vegetables, beverages and other plant based dietary components. In: Crozier A, Clifford MN, Ashihara H, eds. *Plant Secondary Metabolites: Occurrence, Structure and Role in the Human Diet*. Blackwell Publishing Ltd; 2006. pp. 208–302.
- [388] Gülçin İ. Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid). *Toxicology* 2006;217:213–20.
- [389] Yang WS, Jeong D, Yi YS, *et al.* IRAK1/4-targeted anti-inflammatory action of caffeic acid. *Mediators Inflamm* 2013;2013:518183.
- [390] Guo D, Dou D, Ge L, *et al.* A caffeic acid mediated facile synthesis of silver nanoparticles with powerful anti-cancer activity. *Colloids Surf B Biointerfaces* 2015;134:229–34.
- [391] Mirzaei S, Gholami MH, Zabolian A, *et al.* Caffeic acid and its derivatives as potential modulators of oncogenic molecular pathways: new hope in the fight against cancer. *Pharmacol Res* 2021; 171:105759.
- [392] Kabala-Dzik A, Rzepecka-Stojko A, Kubina R, *et al.* Comparison of two components of propolis: caffeic acid (CA) and caffeic acid phenethyl ester (CAPE) induce apoptosis and cell cycle arrest of breast cancer cells MDA-MB-231. *Molecules* 2017;22:1554.
- [393] Dziedzic A, Kubina R, Kabala-Dzik A, *et al.* Induction of cell cycle arrest and apoptotic response of head and neck squamous carcinoma cells (Detroit 562) by caffeic acid and caffeic acid phenethyl ester derivative. *Evid Based Complement Alternat Med* 2017;2017:6793456.

- [394] Rangwala S, Tsai KY. Roles of the immune system in skin cancer. *Br J Dermatol* 2011;165:953–65.
- [395] Hofbauer GF, Bavnick JNB, Euvrard S. Organ transplantation and skin cancer: basic problems and new perspectives. *Exp Dermatol* 2010;19:473–82.
- [396] Bartsch H, Nair J. Chronic inflammation and oxidative stress in the genesis and perpetuation of cancer: role of lipid peroxidation, DNA damage, and repair. *Langenbecks Arch Surg* 2006;391:499–510.
- [397] Zhou Y, Jiang Z, Lu H, *et al.* Recent advances of natural polyphenols activators for Keap1-Nrf2 signalling pathway. *Chem Biodivers* 2019;16:e1900400.
- [398] Scapagnini G, Sonya V, Nader AG, *et al.* Modulation of Nrf2/ARE pathway by food polyphenols: a nutritional neuroprotective strategy for cognitive and neurodegenerative disorders. *Mol Neurobiol* 2011;44:192–201.
- [399] Magesh S, Chen Y, Hu L. Small molecule modulators of Keap1-Nrf2-ARE pathway as potential preventive and therapeutic agents. *Med Res Rev* 2012;32:687–726.
- [400] Yang G, Fu Y, Malakhova M, *et al.* Caffeic acid directly targets ERK1/2 to attenuate solar UV-induced skin carcinogenesis. *Cancer Prev Res (Phila)* 2014;7:1056–66.
- [401] Kudugunti SK, Vad NM, Ekogbo E, *et al.* Efficacy of caffeic acid phenethyl ester (CAPE) in skin B16-F0 melanoma tumor bearing C57BL/6 mice. *Invest New Drugs* 2011;29:52–62.
- [402] Balupillai A, Prasad RN, Ramasamy K, *et al.* Caffeic acid inhibits UVB-induced inflammation and photocarcinogenesis through activation of peroxisome proliferator-activated receptor- γ in mouse skin. *Photochem Photobiol* 2015;91:1458–68.
- [403] Balupillai A, Nagarajan RP, Ramasamy K, *et al.* Caffeic acid prevents UVB radiation induced photocarcinogenesis through regulation of PTEN signalling in human dermal fibroblasts and mouse skin. *Toxicol Appl Pharmacol* 2018;352:87–96.
- [404] Agilan B, Rajendra Prasad N, Kanimozhi G, *et al.* Caffeic acid inhibits chronic UVB-induced cellular proliferation through JAK-STAT 3 signalling in mouse skin. *Photochem Photobiol* 2016;92:467–74.
- [405] Mileo AM, Miccadei S. Polyphenols as modulator of oxidative stress in cancer disease: new therapeutic strategies. *Oxid Med Cell Longev* 2016;2016:6475624.
- [406] Wang SJ, Zeng J, Yang BK, *et al.* Bioavailability of caffeic acid in rats and its absorption properties in the Caco-2 cell model. *Pharm Biol* 2014;52:1150–7.
- [407] D'Archivio M, Filesi C, Di Benedetto R, *et al.* Polyphenols, dietary sources and bioavailability. *Ann Ist Super Sanità* 2007;43:348–61.
- [408] Setty AR, Sigal LH. Herbal medications commonly used in the practice of rheumatology: mechanisms of action, efficacy, and side effects. *Semin Arthr Rheum* 2005;34:773–84.
- [409] Boz H. Ferulic acid in cereals - a review. *Czech J Food Sci* 2015;33:1–7.
- [410] Kähkönen MP, Hopia AI, Vuorela HJ, *et al.* Antioxidant activity of plant extracts containing phenolic compounds. *J Agric Food Chem* 1999;47:3954–62.
- [411] Kumar N, Goel N. Phenolic acids: natural versatile molecules with promising therapeutic applications. *Biotechnol Rep* 2019;24:e00370.
- [412] Zheng W, Wang SY. Antioxidant activity and phenolic compounds in selected herbs. *J Agric Food Chem* 2001;49:5165–70.
- [413] Peterson J, Dwyer J, Adlercreutz H, *et al.* Dietary lignans: physiology and potential for cardiovascular disease risk reduction. *Nutr Rev* 2010;68:571–603.
- [414] Özcan MM, Chalchat JC. Chemical composition and antifungal activity of rosemary (*Rosmarinus officinalis* L.) oil from Turkey. *Int J Food Sci Nutr* 2008;59:691–8.
- [415] Rose DJ, Patterson JA, Hamaker BR. Structural differences among alkali-soluble arabinoxylans from maize (*Zea mays*), rice (*Oryza sativa*), and wheat (*Triticum aestivum*) brans influence human fecal fermentation profiles. *J Agric Food Chem* 2010;58:493–9.
- [416] Srinivasan M, Sudheer AR, Menon VP. Ferulic acid: therapeutic potential through its antioxidant property. *J Clin Biochem Nutr* 2007;40:92–100.
- [417] Lin XF, Min W, Luo D. Anticarcinogenic effect of ferulic acid on ultraviolet-B irradiated human keratinocyte HaCaT cells. *J Med Plants Res* 2010;4:1686–94.
- [418] Rajendran I, Ponrasu T, Rajaram R, *et al.* The apoptotic effect of ferulic acid-synthesized gold nanoparticles against human epidermoid carcinoma (A431) cells via activation of caspase-3 pathway. *J Drug Deliv Sci Technol* 2021;63:102478.
- [419] Khalid EB, Ayman EMEK, Rahman H, *et al.* Natural products against cancer angiogenesis. *Tumor Biol* 2016;37:14513–36.
- [420] Alias LM, Manoharan S, Vellaichamy L, *et al.* Protective effect of ferulic acid on 7,12-dimethylbenz[a]anthracene-induced skin carcinogenesis in Swiss albino mice. *Exp Toxicol Pathol* 2009;61:205–14.
- [421] Choi YS, Sohn YW. Anticancer effect of ferulic acid on cultured human skin melanoma cells. *J Exp Biomed Sci* 2006;12:457–61.
- [422] Murakami A, Nakamura Y, Koshimizu K, *et al.* FA15, a hydrophobic derivative of ferulic acid, suppresses inflammatory responses and skin tumor promotion: comparison with ferulic acid. *Cancer Lett* 2002;180:21–29.
- [423] Diaconeasa Z, Stirbu I, Xiao J, *et al.* Anthocyanins, vibrant color pigments, and their role in skin cancer prevention. *Biomedicines* 2020;8:336.
- [424] Korać RR, Khambolja KM. Potential of herbs in skin protection from ultraviolet radiation. *Pharmacogn Rev* 2011;5:164.
- [425] Hamid AA, Aiyelaagbe OO, Usman LA, *et al.* Antioxidants: its medicinal and pharmacological applications. *Afr J Pure Appl Chem* 2010;4:142–51.
- [426] Offord EA, Gautier JC, Avanti O, *et al.* Photoprotective potential of lycopene, β -carotene, vitamin E, vitamin C and carnosic acid in UVA-irradiated human skin fibroblasts. *Free Radic Biol Med* 2002;32:1293–303.
- [427] Pullar JM, Carr AC, Vissers MC. The roles of vitamin C in skin health. *Nutrients* 2017;9:866.
- [428] Grune T, Lietz G, Palou A, *et al.* β -Carotene is an important vitamin A source for humans. *J Nutr* 2010;140:2268S–85S.
- [429] Amao I. Health benefits of fruits and vegetables: review from Sub-Saharan Africa. In: Asaduzzaman M, Asao T, eds. *Vegetables: Importance of Quality Vegetables to Human Health*. Springer Nature; 2018. pp. 33–53.
- [430] Bulló M, Casas-Agustench P, Amigó-Corregi P, *et al.* Inflammation, obesity and comorbidities: the role of diet. *Public Health Nutr* 2007;10(10A):1164–72.
- [431] Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002;420:860–7.
- [432] Rolls BJ, Drewnowski A, Ledikwe JH. Changing the energy density of the diet as a strategy for weight management. *J Am Diet Assoc* 2005;105:98–103.
- [433] Karimi K, Lindgren TH, Koch CA, *et al.* Obesity as a risk factor for malignant melanoma and non-melanoma skin cancer. *Rev Endocr Metab Dis* 2016;17:389–403.
- [434] Williams Merten J, King JL, Walsh-Childers K, *et al.* Skin cancer risk and other health risk behaviors: a scoping review. *Am J Lifestyle Med* 2016;11:182–96.
- [435] Iqbal J, Abbasi BA, Ahmad R, *et al.* Potential phytochemicals in the fight against skin cancer: current landscape and future perspectives. *Biomed Pharmacother* 2019;109:1381–93.
- [436] Chamcheu JC, Roy T, Uddin MB, *et al.* Role and therapeutic targeting of the PI3K/Akt/mTOR signalling pathway in skin cancer: a review of current status and future trends on natural and synthetic agents therapy. *Cells* 2019;8:803.
- [437] Athar M, Tang X, Lee JL, *et al.* Hedgehog signalling in skin development and cancer. *Exp Dermatol* 2006;15:667–77.
- [438] George BP, Chandran R, Abrahamse H. Role of phytochemicals in cancer chemoprevention: insights. *Antioxidants* 2021;10:1455.
- [439] Cháirez-Ramírez MH, de la Cruz-López KG, García-Carrancá A. Polyphenols as antitumor agents targeting key players in cancer-driving signalling pathways. *Front Pharmacol* 2021;12:710304.
- [440] Pandey P, Khan F. A mechanistic review of the anticancer potential of hesperidin, a natural flavonoid from citrus fruits. *Nutr Res* 2021;92:21–31.
- [441] Choi J, Lee DH, Jang H, *et al.* Naringenin exerts anticancer effects by inducing tumor cell death and inhibiting angiogenesis in malignant melanoma. *Int J Med Sci* 2020;17:3049.
- [442] Mintie CA, Singh CK, Ahmad N. Whole fruit phytochemicals combating skin damage and carcinogenesis. *Transl Oncol* 2020;13:146–56.
- [443] Neagu M, Constantin C, Caruntu C, *et al.* Inflammation: a key process in skin tumorigenesis. *Oncol Lett* 2019;17:4068–84.
- [444] Li Y, Yao J, Han C, *et al.* Quercetin, inflammation and immunity. *Nutrients* 2016;8:167.
- [445] Al Robaee AA. Awareness to sun exposure and use of sunscreen by the general population. *Bosn J Basic Med Sci* 2010;10:314–8.

- [446] Chadni M, Isidore E, Lagalle F, *et al.* Optimization of the supercritical extraction of rosmarinic acid from clary sage residue and the antioxidant activity of the extracts. *J Supercrit Fluids* 2023;193:105830.
- [447] Erkan N, Ayranci G, Ayranci E. Antioxidant activities of rosemary (*Rosmarinus officinalis* L.) extract, blackseed (*Nigella sativa* L.) essential oil, carnosic acid, rosmarinic acid and sesamol. *Food Chem* 2008;110:76–82.
- [448] Wang H, Provan GJ, Helliwell K. Determination of rosmarinic acid and caffeic acid in aromatic herbs by HPLC. *Food Chem* 2004;87:307–11.
- [449] Kiferle C, Lucchesini M, Mensuali-Sodi A, *et al.* Rosmarinic acid content in basil plants grown in vitro and in hydroponics. *Open Life Sci* 2011;6:946–57.
- [450] Perry PL, Shetty K. A model for involvement of proline during pseudomonas-mediated stimulation of rosmarinic acid levels in oregano shoot clones. *Food Biotechnol* 1999;13:137–54.
- [451] Ashoush YAM, Ali AMF, Abozid MM, *et al.* Comparative study between celery leaves and broccoli flowers for their chemical composition and amino acids as well as phenolic and flavonoid compounds. *Menoufia Jf Agric Biotechnol* 2017;2:1–13.
- [452] Gomes PG, Veloso AF, Maynard IF, *et al.* Integrative process to extract chlorophyll and purify rosmarinic acid from rosemary leaves (*Rosmarinus officinalis*). *J Chem Technol Biotechnol* 2020;95:1503–10.
- [453] Carnat AP, Carnat A, Fraisse D, *et al.* The aromatic and polyphenolic composition of lemon balm (*Melissa officinalis* L. subsp. *officinalis*) tea. *Pharm Acta Helv* 1998;72:301–5.
- [454] David IG, Buleandră M, Popa DE, *et al.* Voltammetric determination of polyphenolic content as rosmarinic acid equivalent in tea samples using pencil graphite electrodes. *J Food Sci Technol* 2016;53:2589–96.
- [455] Andrade P, Ferreres F, Gil MI, *et al.* Determination of phenolic compounds in honeys with different floral origin by capillary zone electrophoresis. *Food Chem* 1997;60:79–84.
- [456] Mandal SK, Das A, Dey S, *et al.* Bioactivities of allicin and related organosulfur compounds from garlic: overview of the literature since 2010. *Egypt J Chem* 62 (Special Issue - Part 1 Innovat Chem) 2019;62:1–11.
- [457] Prati P, Henrique CM, Souza ASD, *et al.* Evaluation of allicin stability in processed garlic of different cultivars. *Food Sci Technol* 2014;34:623–8.
- [458] Yin MC, Cheng WS. Antioxidant activity of several *Allium* members. *J Agric Food Chem* 1998;46:4097–101.
- [459] Melouk SA, Hassan MA, Elwan MWM, *et al.* Horticultural, chemical and genetic diversity using SSR markers in Leek germplasm collection. *Sci Hortic* 2023;311:111782.
- [460] Ghodrati Azadi H, Ghaffari SM, Riazi GH, *et al.* Antiproliferative activity of chloroformic extract of Persian Shallot, *Allium hirtifolium*, on tumor cell lines. *Cytotechnology* 2008;56:179–85.
- [461] Gu C, Ma H, Tuly JA, *et al.* Effects of catalytic infrared drying in combination with hot air drying and freeze drying on the drying characteristics and product quality of chives. *LWT* 2022;161:113363.
- [462] de Figueiredo SM, Binda NS, Nogueira-Machado JA, *et al.* The antioxidant properties of organosulfur compounds (sulforaphane). *Recent Pat Endocr Metab Immune Drug Discov* 2015;9:24–39.
- [463] Moon JK, Kim JR, Ahn YJ, *et al.* Analysis and anti-Helicobacter activity of sulforaphane and related compounds present in broccoli (*Brassica oleracea* L.) sprouts. *J Agric Food Chem* 2010;58:6672–7.
- [464] Čekey N, Šlosár M, Uher A, *et al.* The effect of nitrogen and sulphur fertilization on the yield and content of sulforaphane and nitrates in cauliflower. *Acta Univ Agric Silvicae Mendel Brun* 2011;59:17–22.
- [465] Bryant CS, Kumar S, Chamala S, *et al.* Sulforaphane induces cell cycle arrest by protecting RB-E2F-1 complex in epithelial ovarian cancer cells. *Mol Cancer* 2010;9:47.
- [466] Sasaki K, Neyazaki M, Shindo K, *et al.* Quantitative profiling of glucosinolates by LC–MS analysis reveals several cultivars of cabbage and kale as promising sources of sulforaphane. *J Chromatogr B Anal Technol Biomed Life Sci* 2012;903:171–6.
- [467] Kim J. Pre-clinical neuroprotective evidences and plausible mechanisms of sulforaphane in Alzheimer's disease. *Int J Mol Sci* 2021;22:2929.
- [468] Herr I, Lozanovski V, Houben P, *et al.* Sulforaphane and related mustard oils in focus of cancer prevention and therapy. *Wien Med Wochenschr* 2013;163:80–8.
- [469] Dosz EB, Jeffery EH. Modifying the processing and handling of frozen broccoli for increased sulforaphane formation. *J Food Sci* 2013;78:H1459–63.
- [470] Belyakova T, Morozova O, Antonceva E, *et al.* Perspective processing of turnip root (*Brassica rapa*) as a source of sulforafan. In: *E3S Web of Conferences*, Vol. 164. EDP Sciences; 2020. 06028. 6pp.
- [471] Rosillo MA, Sanchez-Hidalgo M, Cárdeno A, *et al.* Protective effect of ellagic acid, a natural polyphenolic compound, in a murine model of Crohn's disease. *Biochem Pharmacol* 2011;82:737–45.
- [472] Zafrilla P, Ferreres F, Tomás-Barberán FA. Effect of processing and storage on the antioxidant ellagic acid derivatives and flavonoids of red raspberry (*Rubus idaeus*) jams. *J Agric Food Chem* 2001;49:3651–5.
- [473] Đurić M, Mašković P, Murtić S, *et al.* Quantitation of ellagic acid in blackberries. *Hem Ind* 2014;68:241–5.
- [474] Usta C, Ozdemir S, Schiariti M, *et al.* The pharmacological use of ellagic acid-rich pomegranate fruit. *Int J Food Sci Nutr* 2013;64:907–13.
- [475] Vattem DA, Shetty K. Ellagic acid production and phenolic antioxidant activity in cranberry pomace (*Vaccinium macrocarpon*) mediated by *Lentinus edodes* using a solid-state system. *Process Biochem* 2003;39:367–79.
- [476] Flores-Cordova M, Muñoz-Márquez E, Muñoz-Márquez E, *et al.* Phytochemical composition and antioxidant capacity in Mexican pecan nut. *Emir J Food Agric* 2017;29:346–50.
- [477] Han DH, Lee MJ, Kim JH. Antioxidant and apoptosis-inducing activities of ellagic acid. *Anticancer Res* 2006;26(5A):3601–6.
- [478] Weisburg JH, Schuck AG, Reiss SE, *et al.* Ellagic acid, a dietary polyphenol, selectively cytotoxic to HSC-2 oral carcinoma cells. *Anticancer Res* 2013;33:1829–36.
- [479] de Melo CL, Queiroz MGR, Arruda Filho ACV, *et al.* Betulinic acid, a natural pentacyclic triterpenoid, prevents abdominal fat accumulation in mice fed a high-fat diet. *J Agric Food Chem* 2009;57:8776–81.
- [480] Alhazmi H. Extraction of phytochemicals betulin and betulinic acid from the chaga mushroom and their effect on MCF-7 cells [Doctoral dissertation], 2018.
- [481] Hossain U, Das AK, Ghosh S, *et al.* An overview on the role of bioactive α -glucosidase inhibitors in ameliorating diabetic complications. *Food Chem Toxicol* 2020;145:111738.
- [482] Liu J, Chen P, Yao W, *et al.* Subcritical water extraction of betulinic acid from birch bark. *Ind Crops Prod* 2015;74:557–65.
- [483] Madunić J, Madunić IV, Gajski G, *et al.* Apigenin: a dietary flavonoid with diverse anticancer properties. *Cancer Lett* 2018;413:11–22.
- [484] Poureini F, Najafpour GD, Nikzad M, *et al.* Loading of apigenin extracted from parsley leaves on colloidal core-shell nanocomposite for bioavailability enhancement. *Colloids Surf A Physicochem Eng Aspects* 2021;625:126867.
- [485] Srivastava JK, Shankar E, Gupta S. Chamomile: a herbal medicine of the past with bright future. *Mol Med Rep* 2010;3:895–901.
- [486] Kruma Z, Andjelkovic M, Verhe R, *et al.* Phenolic compounds in basil, oregano and thyme. *Foodbalt* 2008;5:99–103.
- [487] Oyenihni OR, Oyenihni AB, Alabi TD, *et al.* Reactive oxygen species: key players in the anticancer effects of apigenin? *J Food Biochem* 2022;46:e14060.
- [488] Namsi A, Nury T, Hamdouni H, *et al.* Induction of neuronal differentiation of murine N2a cells by two polyphenols present in the Mediterranean diet mimicking neurotrophins activities: resveratrol and apigenin. *Diseases* 2018;6:67.
- [489] Patel D, Shukla S, Gupta S. Apigenin and cancer chemoprevention: progress, potential and promise. *Int J Oncol* 2007;30:233–45.
- [490] Pandino G, Courts FL, Lombardo S, *et al.* Caffeoylquinic acids and flavonoids in the immature inflorescence of globe artichoke, wild cardoon, and cultivated cardoon. *J Agric Food Chem* 2010;58:1026–31.
- [491] Pająk P, Socha R, Galkowska D, *et al.* Phenolic profile and antioxidant activity in selected seeds and sprouts. *Food Chem* 2014;143:300–6.
- [492] Meckelmann SW, Riegel DW, Van Zonneveld MJ, *et al.* Compositional characterization of native *Peruvian chili* peppers (*Capsicum* spp.). *J Agric Food Chem* 2013;61:2530–7.
- [493] Rowell KM, Beisel CG. Isolation of gram quantities of a rhamnoglucoside of apigenin from grapefruit. *J Food Sci* 1963;28:195–7.
- [494] Szczepaniak O, Ligaj M, Stuper-Szablewska K, *et al.* Genoprotective effect of cornelian cherry (*Cornus mas* L.) phytochemicals, electrochemical and ab initio interaction study. *Biomed Pharmacother* 2022;152:113216.
- [495] McKay DL, Blumberg JB. A review of the bioactivity and potential health benefits of chamomile tea (*Matricaria recutita* L.). *Phytother Res* 2006;20:519–30.

- [496] Pais JM, Pereira B, Paz FAA, *et al.* Solid γ -cyclodextrin inclusion compound with gingerols, a multi-component guest: preparation, properties and application in yogurt. *Biomolecules* 2020;10:344.
- [497] Schwertner HA, Rios DC, Pascoe JE. Variation in concentration and labeling of ginger root dietary supplements. *Obstet Gynecol* 2006;107:1337–43.
- [498] David AVA, Arulmoli R, Parasuraman S. Overviews of biological importance of quercetin: a bioactive flavonoid. *Pharmacogn Rev* 2016;10:84.
- [499] Hollman PC, Van Trijp JM, Buysman MN, *et al.* Relative bioavailability of the antioxidant flavonoid quercetin from various foods in man. *FEBS Lett* 1997;418:152–6.
- [500] Wilms LC, Hollman PC, Boots AW, *et al.* Protection by quercetin and quercetin-rich fruit juice against induction of oxidative DNA damage and formation of BPDE-DNA adducts in human lymphocytes. *Mutat Res* 2005;582:155–62.
- [501] Neto CC, Dao CA, Salvas MR, *et al.* Variation in concentration of phenolic acid derivatives and quercetin glycosides in foliage of cranberry that may play a role in pest deterrence. *J Am Soc Hortic Sci* 2010;135:494–500.
- [502] Schmitzer V, Veberic R, Stampar F. European elderberry (*Sambucus nigra* L.) and American Elderberry (*Sambucus canadensis* L.): botanical, chemical and health properties of flowers, berries and their products. *Berries Prop Consum Nutr* 2012;2012:127–44.
- [503] Snyder SM, Zhao B, Luo T, *et al.* Consumption of quercetin and quercetin-containing apple and cherry extracts affects blood glucose concentration, hepatic metabolism, and gene expression patterns in obese C57BL/6J high fat-fed mice. *J Nutr* 2016;146:1001–7.
- [504] Eftekhari M, Alizadeh M, Ebrahimi P. Evaluation of the total phenolics and quercetin content of foliage in mycorrhizal grape (*Vitis vinifera* L.) varieties and effect of postharvest drying on quercetin yield. *Ind Crop Prod* 2012;38:160–5.
- [505] Careri M, Elviri L, Mangia A, *et al.* Spectrophotometric and coulometric detection in the high-performance liquid chromatography of flavonoids and optimization of sample treatment for the determination of quercetin in orange juice. *J Chromatogr A* 2000;881:449–60.
- [506] Farag RS, Abdel-Latif MS, Emam SS, *et al.* Phytochemical screening and polyphenol constituents of pomegranate peels and leave juices. *Agric Soil Sci* 2014;1:86–93.
- [507] McAnlis GT, McEneny J, Pearce J, *et al.* Absorption and antioxidant effects of quercetin from onions, in man. *Eur J Clin Nutr* 1999;53:92–6.
- [508] Wiczowski W, Romaszko J, Bucinski A, *et al.* Quercetin from shallots (*Allium cepa* L. var. *aggregatum*) is more bioavailable than its glucosides. *J Nutr* 2008;138:885–8.
- [509] Yang I, Jayaprakasha GK, Patil B. In vitro digestion with bile acids enhances the bioaccessibility of kale polyphenols. *Food Funct* 2018;9:1235–44.
- [510] Zein N, Yassin F, Othman M. Quercetin extracted from broccoli attenuates the renewal of hepatic cells via downregulation of TGF β -1 and arresting of HSCs activation in mice. *Biointerface Res Appl Chem* 2021;11:12348–63.
- [511] Bozack A. Effect of organic and conventional farming practices on quercetin content in spinach, *Spinacia oleracea*. *Quercetin Content Spinach* 2006;5:1–15.
- [512] Hirose Y, Fujita T, Ishii T, *et al.* Antioxidative properties and flavonoid composition of *Chenopodium quinoa* seeds cultivated in Japan. *Food Chem* 2010;119:1300–6.
- [513] Zhao G, Zou L, Wang Z, *et al.* Pharmacokinetic profile of total quercetin after single oral dose of tartary buckwheat extracts in rats. *J Agric Food Chem* 2011;59:4435–41.
- [514] Kim IS, Yang MR, Lee OH, *et al.* Antioxidant activities of hot water extracts from various spices. *Int J Mol Sci* 2011;12:4120–31.
- [515] Jin D, Hakamata H, Takahashi K, *et al.* Determination of quercetin in human plasma after ingestion of commercial canned green tea by semi-micro HPLC with electrochemical detection. *Biomed Chromatogr* 2004;18:662–6.
- [516] McDonald MS, Hughes M, Burns J, *et al.* Survey of the free and conjugated myricetin and quercetin content of red wines of different geographical origins. *J Agric Food Chem* 1998;46:368–75.
- [517] Guo AJ, Choi RC, Zheng KY, *et al.* Kaempferol as a flavonoid induces osteoblastic differentiation via estrogen receptor signalling. *Chin Med* 2012;7:10.
- [518] Dehkharghanian M, Adenier H, Vijayalakshmi MA. Study of flavonoids in aqueous spinach extract using positive electrospray ionisation tandem quadrupole mass spectrometry. *Food Chem* 2010;121:863–70.
- [519] Gliszczynska-Swiglo A, Kaluzewicz A, Lemańska K, *et al.* The effect of solar radiation on the flavonol content in broccoli inflorescence. *Food Chem* 2007;100:241–5.
- [520] Nirmala P, Ramanathan M. Effect of kaempferol on lipid peroxidation and antioxidant status in 1,2-dimethyl hydrazine induced colorectal carcinoma in rats. *Eur J Pharmacol* 2011;654:75–9.
- [521] Nielsen JK, Nørbæk R, Olsen CE. Kaempferol tetraglucosides from cabbage leaves. *Phytochemistry* 1998;49:2171–6.
- [522] Castillo-Muñoz N, Gómez-Alonso S, García-Romero E, *et al.* Flavonol profiles of Vitis vinifera red grapes and their single-cultivar wines. *J Agric Food Chem* 2007;55:992–1002.
- [523] Cho HJ, Park JHY. Kaempferol induces cell cycle arrest in HT-29 human colon cancer cells. *J Cancer Prev* 2013;18:257.
- [524] Karabagias IK, Dimitriou E, Kontakos S, *et al.* Phenolic profile, colour intensity, and radical scavenging activity of Greek unifloral honeys. *Eur Food Res Technol* 2016;242:1201–10.
- [525] Park JS, Rho HS, Kim DH, *et al.* Enzymatic preparation of kaempferol from green tea seed and its antioxidant activity. *J Agric Food Chem* 2006;54:2951–6.
- [526] Wang H, Helliwell K. Determination of flavonols in green and black tea leaves and green tea infusions by high-performance liquid chromatography. *Food Res Int* 2001;34:223–7.
- [527] Singh A, Jain A, Sarma BK, *et al.* Rhizosphere competent microbial consortium mediates rapid changes in phenolic profiles in chickpea during *Sclerotium rolfsii* infection. *Microbiol Res* 2014;169:353–60.
- [528] Manoj BS, Gupta M, Jeelani MI, *et al.* Metabolic footprints of chitosan primed red kidney bean under restricted irrigation. *Int J Biol Macromol* 2022;208:367–80.
- [529] Frémont L. Biological effects of resveratrol. *Life Sci* 2000;66:663–73.
- [530] Leyva-Porras C, Saavedra-Leos MZ, Cervantes-González E, *et al.* Spray drying of blueberry juice-maltodextrin mixtures: evaluation of processing conditions on content of resveratrol. *Antioxidants* 2019;8:437.
- [531] Rocha-González HI, Ambriz-Tututi M, Granados-Soto V. Resveratrol: a natural compound with pharmacological potential in neurodegenerative diseases. *CNS Neurosci Ther* 2008;14:234–47.
- [532] Counet C, Callemien D, Collin S. Chocolate and cocoa: new sources of *trans*-resveratrol and *trans*-piceid. *Food Chem* 2006;98:649–57.
- [533] Ragab AS, Van Fleet J, Jankowski B, *et al.* Detection and quantitation of resveratrol in tomato fruit (*Lycopersicon esculentum* Mill.). *J Agric Food Chem* 2006;54:7175–9.
- [534] Shankar S, Srivastava RK. Bax and Bak genes are essential for maximum apoptotic response by curcumin, a polyphenolic compound and cancer chemopreventive agent derived from turmeric, *Curcuma longa*. *Carcinogenesis* 2007;28:1277–86.
- [535] Weiss DJ, Anderton CR. Determination of catechins in matcha green tea by micellar electrokinetic chromatography. *J Chromatogr A* 2003;1011:173–80.
- [536] Sanlier N, Atik İ, Atik A. A minireview of effects of white tea consumption on diseases. *Trends Food Sci Technol* 2018;82:82–8.
- [537] Oliveira A, Almeida DP, Pintado M. Changes in phenolic compounds during storage of pasteurized strawberry. *Food Bioprocess Technol* 2014;7:1840–6.
- [538] Pavlović AV, Papetti A, Zagorac ĐČD, *et al.* Phenolics composition of leaf extracts of raspberry and blackberry cultivars grown in Serbia. *Ind Crops Prod* 2016;87:304–14.
- [539] Varo MÁ, Martín-Gómez J, Mérida J, *et al.* Bioactive compounds and antioxidant activity of highbush blueberry (*Vaccinium corymbosum*) grown in southern Spain. *Eur Food Res Technol* 2021;247:1199–208.
- [540] Souquet JM, Cheynier V, Brossaud F, *et al.* Polymeric proanthocyanidins from grape skins. *Phytochemistry* 1996;43:509–12.
- [541] Casanova E, Salvadó J, Crescenti A, *et al.* Epigallocatechin gallate modulates muscle homeostasis in type 2 diabetes and obesity by targeting energetic and redox pathways: a narrative review. *Int J Mol Sci* 2019;20:532.
- [542] Yoo KS, Bang H, Pike L, *et al.* Comparing carotene, anthocyanins, and terpenoid concentrations in selected carrot lines of different colors. *Hortic Environ Biotechnol* 2020;61:385–93.
- [543] Wu X, Sun C, Yang L, *et al.* β -Carotene content in sweet potato varieties from China and the effect of preparation on β -carotene retention in the Yanshu No. 5. *Innov Food Sci Emerg Technol* 2008;9:581–6.

- [544] Koh SH, Loh SP. In vitro bioaccessibility of β -carotene in pumpkin and butternut squash subjected to different cooking methods. *Int Food Res J* 2018;25:188–95.
- [545] Tsao R, Yang R. Lutein in selected Canadian crops and agri-food processing by-products and purification by high-speed counter-current chromatography. *J Chromatogr A* 2006;1112:202–8.
- [546] Rock CL, Lovatlo JL, Emenhiser C, et al. Bioavailability of β -carotene is lower in raw than in processed carrots and spinach in women. *J Nutr* 1998;128:913–6.
- [547] Chen X, Wu J, Zhou S, et al. Application of near-infrared reflectance spectroscopy to evaluate the lutein and β -carotene in Chinese kale. *J Food Compos Anal* 2009;22:148–53.
- [548] Ren Y, Bang H, Lee EJ, et al. Levels of phytoene and β -carotene in transgenic honeydew melon (*Cucumis melo L. inodorus*). *Plant Cell Tissue Organ Cult (PCTOC)* 2013;113:291–301.
- [549] Ornelas-Paz JDJ, Failla ML, Yahia EM, et al. Impact of the stage of ripening and dietary fat on in vitro bioaccessibility of β -carotene in 'Ataulfo' mango. *J Agric Food Chem* 2008;56:1511–6.
- [550] Karabulut I, Topcu A, Duran A, et al. Effect of hot air drying and sun drying on color values and β -carotene content of apricot (*Prunus armenica* L.). *LWT-Food Sci Technol* 2007;40:753–8.
- [551] Espíndola KMM, Ferreira RG, Narvaez LEM, et al. Chemical and pharmacological aspects of caffeic acid and its activity in hepatocarcinoma. *Front Oncol* 2019;9:541.
- [552] Reinders RD, Biesterveld S, Bijker PG. Survival of *Escherichia coli* O157: H7 ATCC 43895 in a model apple juice medium with different concentrations of proline and caffeic acid. *Appl Environ Microbiol* 2001;67:2863–6.
- [553] Öztürk A, Demirsoy L, Demirsoy H, et al. Phenolic compounds and chemical characteristics of pears (*Pyrus Communis* L.). *Int J Food Prop* 2015;18:536–46.
- [554] Sellappan S, Akoh CC, Krewer G. Phenolic compounds and antioxidant capacity of Georgia-grown blueberries and blackberries. *J Agric Food Chem* 2002;50:2432–8.
- [555] Levaj B, Dragović-Uzelac V, Delonga K, et al. Polyphenols and volatiles in fruits of two sour cherry cultivars, some berry fruits and their jams. *Food Technol Biotechnol* 2010;48:538–47.
- [556] Nia NN, Hadjmohammadi MR. Amino acids-based hydrophobic natural deep eutectic solvents as a green acceptor phase in two-phase hollow fiber-liquid microextraction for the determination of caffeic acid in coffee, green tea, and tomato samples. *Microchem J* 2021;164:06021.
- [557] Singletary K. Oregano: overview of the literature on health benefits. *Nutr Today* 2010;45:129–38.
- [558] Peng Y, Yuan J, Liu F, et al. Determination of active components in rosemary by capillary electrophoresis with electrochemical detection. *J Pharm Biomed Anal* 2005;39:431–7.
- [559] Areias F, Valente P, Andrade PB, et al. Flavonoids and phenolic acids of sage: influence of some agricultural factors. *J Agric Food Chem* 2000;48:6081–4.
- [560] Banjanin T, Nikolic D, Uslu N, et al. Physicochemical properties, fatty acids, phenolic compounds, and mineral contents of 12 Serbia regional and commercial almond cultivars. *J Food Process Preserv* 2021;45:e15015.
- [561] Pedrosa MM, Muzquiz M, García-Vallejo C, et al. Determination of caffeic and chlorogenic acids and their derivatives in different sunflower seeds. *J Sci Food Agric* 2000;80:459–64.
- [562] Li M, Luo J, Nawaz MA, et al. Phytochemistry, bioaccessibility, and bioactivities of sesame seeds: an overview. *Food Rev Int* 2023;40:309–35.
- [563] Ferri M, Happel A, Zanaroli G, et al. Advances in combined enzymatic extraction of ferulic acid from wheat bran. *New Biotechnol* 2020;56:38–45.
- [564] Liu RH. Whole grain phytochemicals and health. *J Cereal Sci* 2007;46:207–19.
- [565] Kelebek H, Selli S, Canbas A, et al. HPLC determination of organic acids, sugars, phenolic compositions and antioxidant capacity of orange juice and orange wine made from a Turkish cv. Kozan. *Microchem J* 2009;91:187–92.
- [566] Lee J, Chan BLS, Mitchell AE. Identification/quantification of free and bound phenolic acids in peel and pulp of apples (*Malus domestica*) using high resolution mass spectrometry (HRMS). *Food Chem* 2017;215:301–10.
- [567] Smith BG, Harris PJ. Ferulic acid is esterified to glucuronarabinoxylans in pineapple cell walls. *Phytochemistry* 2001;56:513–9.
- [568] Bursal E, Köksal E, Gülçin İ, et al. Antioxidant activity and polyphenol content of cherry stem (*Cerasus avium* L.) determined by LC–MS/MS. *Food Res Int* 2013;51:66–74.
- [569] Lichtenthaler HK, Schweiger J. Cell wall bound ferulic acid, the major substance of the blue-green fluorescence emission of plants. *J Plant Physiol* 1998;152:272–82.
- [570] Rodríguez García SL, Raghavan V. Microwave-assisted extraction of phenolic compounds from broccoli (*Brassica oleracea*) stems, leaves, and florets: optimization, characterization, and comparison with maceration extraction. *Rec Progr Nutr* 2022;2:1–20.
- [571] Szczepańska J, Barba FJ, Skapska S, et al. High pressure processing of carrot juice: effect of static and multi-pulsed pressure on the polyphenolic profile, oxidoreductases activity and colour. *Food Chem* 2020;307:125549.
- [572] Min JY, Kang SM, Park DJ, et al. Enzymatic release of ferulic acid from *Ipomoea batatas* L. (sweet potato) stem. *Biotechnol Bioprocess Eng* 2006;11:372–6.
- [573] Vallverdú-Queralt A, Regueiro J, Martínez-Huélamo M, et al. A comprehensive study on the phenolic profile of widely used culinary herbs and spices: rosemary, thyme, oregano, cinnamon, cumin and bay. *Food Chem* 2014;154:299–307.
- [574] Eliasson C, Kamal-Eldin A, Andersson R, et al. High-performance liquid chromatographic analysis of secoisolaricresinol diglucoside and hydroxycinnamic acid glucosides in flaxseed by alkaline extraction. *J Chromatogr A* 2003;1012:151–9.
- [575] Khezeli T, Daneshfar A, Sahraei R. A green ultrasonic-assisted liquid-liquid microextraction based on deep eutectic solvent for the HPLC-UV determination of ferulic, caffeic and cinnamic acid from olive, almond, sesame and cinnamon oil. *Talanta* 2016;150:577–85.
- [576] Chen L, Liu Y, Liu H, et al. Identification and expression analysis of MATE genes involved in flavonoid transport in blueberry plants. *PLoS One* 2015;10:e0118578.
- [577] Die JV, Jones RW, Ogden EL, et al. Characterization and analysis of anthocyanin-related genes in wild-type blueberry and the pink-fruited mutant cultivar 'Pink Lemonade': new insights into anthocyanin biosynthesis. *Agronomy* 2020;10:1296.
- [578] Kähkönen MP, Heinonen M. Antioxidant activity of anthocyanins and their aglycons. *J Agric Food Chem* 2003;51:628–33.
- [579] Lee YM, Yoon Y, Yoon H, et al. Dietary anthocyanins against obesity and inflammation. *Nutrients* 2017;9:1089.
- [580] Cassidy A. Berry anthocyanin intake and cardiovascular health. *Mol Aspects Med* 2018;61:76–82.
- [581] Wang D, Guo H, Yang H, et al. Pterostilbene, an active constituent of blueberries, suppresses proliferation potential of human cholangiocarcinoma via enhancing the autophagic flux. *Front Pharmacol* 2019;10:1238.
- [582] Campelo APBS. The use of pterostilbene in the treatment of skin cancer: a literature review. *Clin Surg* 2022;8:1–4.
- [583] Shivemba A, Ojinnaka D. Determination of vitamin C and total phenolic in fresh and freeze dried blueberries and the antioxidant capacity of their extracts. *Integr Food Nutr Metab* 2017;4:1–5.
- [584] Maya-Cano DA, Arango-Varela S, Santa-Gonzalez GA. Phenolic compounds of blueberries (*Vaccinium* spp.) as a protective strategy against skin cell damage induced by ROS: a review of antioxidant potential and antiproliferative capacity. *Heliyon* 2021;7:e06297.
- [585] Panche AN, Diwan AD, Chandra SR. Flavonoids: an overview. *J Nutr Sci* 2016;5:e47.
- [586] Coates EM, Popa G, Gill CI, et al. Colon-available raspberry polyphenols exhibit anti-cancer effects on in vitro models of colon cancer. *J Carcinog* 2007;6:4.
- [587] Katta R, Brown DN. Diet and skin cancer: the potential role of dietary antioxidants in non-melanoma skin cancer prevention. *J Skin Cancer* 2015;2015:893149.
- [588] Duncan FJ, Martin JR, Wulff BC, et al. Topical treatment with black raspberry extract reduces cutaneous UVB-induced carcinogenesis and inflammation. *Cancer Prev Res (Phila)* 2009;2:665–72.
- [589] Oberyszyn T. Effects of black raspberries on UV-induced cutaneous inflammation and tumor development. In: Seeram NP, Stoner GD, eds. *Berries and Cancer Prevention*. Springer; 2011. pp. 131–42.
- [590] Mace TA, King SA, Ameen Z, et al. Bioactive compounds or metabolites from black raspberries modulate T lymphocyte proliferation, myeloid cell differentiation and Jak/STAT signalling. *Cancer Immunol Immunother* 2014;63:889–900.

- [591] Satpal D, Kaur J, Bhadariya V, *et al.* Actinidia deliciosa (Kiwi fruit): a comprehensive review on the nutritional composition, health benefits, traditional utilization, and commercialization. *J Food Process Preserv* 2021;45:e15588.
- [592] Kim YM, Park YS, Park YK, *et al.* Characterization of bioactive ligands with antioxidant properties of kiwifruit and persimmon cultivars using in vitro and in silico studies. *Appl Sci* 2020;10:4218.
- [593] Azeem M, Hanif M, Mahmood K, *et al.* An insight into anticancer, antioxidant, antimicrobial, antidiabetic and anti-inflammatory effects of quercetin: a review. *Polym Bull* 2023;80:241–62.
- [594] Wang J, Fang X, Ge L, *et al.* Antitumor, antioxidant and anti-inflammatory activities of kaempferol and its corresponding glycosides and the enzymatic preparation of kaempferol. *PLoS One* 2018;13:e0197563.
- [595] Ampomah-Dwamena C, McGhie T, Wibisono R, *et al.* The kiwifruit lycopene beta-cyclase plays a significant role in carotenoid accumulation in fruit. *J Exp Bot* 2009;60:3765–79.
- [596] Nishiyama I, Fukuda T, Oota T. Cultivar difference in chlorophyll, lutein and β -carotene content in the fruit of kiwifruit and other *Actinidia* species. *Acta Horticult* 2006;753:473–8.
- [597] Sommerburg O, Keunen JE, Bird AC, *et al.* Fruits and vegetables that are sources for lutein and zeaxanthin: the macular pigment in human eyes. *Br J Ophthalmol* 1998;82:907–10.
- [598] Chauhan M, Garg V, Zia G, *et al.* Potential role of phytochemicals of fruits and vegetables in human diet. *Res J Pharm Technol* 2020;13:1587–91.
- [599] El-Raey MA, Ibrahim GE, Eldahshan OA. Lycopene and Lutein; A review for their chemistry and medicinal uses. *J Pharmacogn Phytochem* 2013;2:245–54.
- [600] Kim YE, Cho CH, Kang H, *et al.* Kiwifruit of *Actinidia eriantha* cv. Bidan has in vitro antioxidative, anti-inflammatory and immunomodulatory effects on macrophages and splenocytes isolated from male BALB/c mice. *Food Sci Biotechnol* 2018;27:1503–11.
- [601] Kou L, Zhu Z, Redington C, *et al.* Potential use of kiwifruit extract for treatment of melanoma. *Med Oncol* 2021;38:1–7.
- [602] Kou L, Zhu Z, Fajardo E, *et al.* Harnessing the power of kiwifruit for radiosensitization of melanoma. *Anticancer Res* 2021;41:5945–51.
- [603] Maoto MM, Beswa D, Jideani AI. Watermelon as a potential fruit snack. *Int J Food Prop* 2019;22:355–70.
- [604] Kim CH, Park MK, Kim SK, *et al.* Antioxidant capacity and anti-inflammatory activity of lycopene in watermelon. *Int J Food Sci Technol* 2014;49:2083–91.
- [605] van Breemen RB, Pajkovic N. Multitargeted therapy of cancer by lycopene. *Cancer Lett* 2008;269:339–51.
- [606] Manivannan A, Lee ES, Han K, *et al.* Versatile nutraceutical potentials of watermelon - a modest fruit loaded with pharmaceutically valuable phytochemicals. *Molecules* 2020;25:5258.
- [607] Lester G. Melon (*Cucumis melo* L.) fruit nutritional quality and health functionality. *HortTechnology* 1997;7:222–7.
- [608] Itoh T, Fujita S, Koketsu M, *et al.* Citrulluside H and citrulluside T from young watermelon fruit attenuate ultraviolet B radiation-induced matrix metalloproteinase expression through the scavenging of generated reactive oxygen species in human dermal fibroblasts. *Photodermatol Photoimmunol Photomed* 2021;37:386–94.
- [609] Ascenso A, Ribeiro H, Marques HC, *et al.* Chemoprevention of photocarcinogenesis by lycopene. *Exp Dermatol* 2014;23:874–8.
- [610] Mirza B, Croley CR, Ahmad M, *et al.* Mango (*Mangifera indica* L.): a magnificent plant with cancer preventive and anticancer therapeutic potential. *Crit Rev Food Sci Nutr* 2021;61:2125–51.
- [611] Saha S, Sadhukhan P, Sil PC. Mangiferin: a xanthonoid with multi-potent anti-inflammatory potential. *Biofactors* 2016;42:459–74.
- [612] Shi W, Deng J, Tong R, *et al.* Molecular mechanisms underlying mangiferin-induced apoptosis and cell cycle arrest in A549 human lung carcinoma cells. *Mol Med Rep* 2016;13:3423–32.
- [613] Schagen SK, Zampeli VA, Makrantonaki E, *et al.* Discovering the link between nutrition and skin aging. *Dermatoendocrinology* 2012;4:298–307.
- [614] Balić A, Mokos M. Do we utilize our knowledge of the skin protective effects of carotenoids enough? *Antioxidants* 2019;8:259.
- [615] Li J, Malakhova M, Mottamal M, *et al.* Norathyriol suppresses skin cancers induced by solar ultraviolet radiation by targeting ERK kinases. *Cancer Res* 2012;72:260–70.
- [616] Saleem M, Afaq F, Adhami VM, *et al.* Lupeol modulates NF- κ B and PI3K/Akt pathways and inhibits skin cancer in CD-1 mice. *Oncogene* 2004;23:5203–14.
- [617] Liu RH. Potential synergy of phytochemicals in cancer prevention: mechanism of action. *J Nutr* 2004;134:3479S–85S.
- [618] Ahmad T, Cawood M, Iqbal Q, *et al.* Phytochemicals in *Daucus carota* and their health benefits. *Foods* 2019;8:424.
- [619] Hughes DA. Dietary carotenoids and human immune function. *Nutrition* 2001;17:823–7.
- [620] Boelsma E, Hendriks HF, Roza L. Nutritional skin care: health effects of micronutrients and fatty acids. *Am J Clin Nutr* 2001;73:853–64.
- [621] Alfurayhi R, Huang L, Brandt K. Pathways affected by falcarninol-type polyacetylenes and implications for their anti-inflammatory function and potential in cancer chemoprevention. *Foods* 2023;12:1192.
- [622] Cooper DA, Eldridge AL, Peters JC. Dietary carotenoids and certain cancers, heart disease, and age-related macular degeneration: a review of recent research. *Nutr Rev* 1999;57:201–14.
- [623] Shebaby WN, Mroueh MA, Boukamp P, *et al.* Wild carrot pentane-based fractions suppress proliferation of human HaCaT keratinocytes and protect against chemically-induced skin cancer. *BMC Complement Altern Med* 2017;17:36.
- [624] Natarajmurthy SH, Dharmesh SM. Amelioration of skin cancer in mice by β -carotene and phenolics of carrot (*Daucus carota*). *Am J Biopharmacol Biochem Life Sci* 2015;4:1–25.
- [625] Islam SU, Ahmed MB, Ahsan H, *et al.* An update on the role of dietary phytochemicals in human skin cancer: new insights into molecular mechanisms. *Antioxidants (Basel)* 2020;9:916.
- [626] Otto A, Du Plessis J, Wiechers JW. Formulation effects of topical emulsions on transdermal and dermal delivery. *Int J Cosmet Sci* 2009;31:1–19.
- [627] Bitev M, Desalegn T, Demissie TB, *et al.* Pharmacokinetics and drug-likeness of antidiabetic flavonoids: molecular docking and DFT study. *PLoS One* 2021;16:e0260853.
- [628] Puglia C, Lauro MR, Tirendi GG, *et al.* Modern drug delivery strategies applied to natural active compounds. *Expert Opin Drug Deliv* 2017;14:755–68.
- [629] More MP, Pardeshi SR, Pardeshi CV, *et al.* Recent advances in phytochemical-based nano-formulation for drug-resistant cancer. *Med Drug Discov* 2021;10:100082.
- [630] Thakur L, Ghodasra U, Patel N, *et al.* Novel approaches for stability improvement in natural medicines. *Pharmacogn Rev* 2011;5:48.
- [631] Bonifacio BV, da Silva PB, Ramos MADS, *et al.* Nanotechnology-based drug delivery systems and herbal medicines: a review. *Int J Nanomedicine* 2014;9:1–15.